



Recent strategies of increasing metal tolerance and phytoremediation potential using genetic transformation of plants

Aleksandra Koźmińska¹ · Alina Wiszniewska¹ · Ewa Hanus-Fajerska¹ · Ewa Muszyńska²

Received: 25 September 2017 / Accepted: 18 December 2017 / Published online: 3 January 2018
© The Author(s) 2018. This article is an open access publication

Abstract

Avoidance and reduction of soil contamination with heavy metals is one of the most serious global challenges. Nowadays, science offers us new opportunities of utilizing plants to extract toxic elements from the soil by means of phytoremediation. Plant abilities to uptake, translocate, and transform heavy metals, as well as to limit their toxicity, may be significantly enhanced via genetic engineering. This paper provides a comprehensive review of recent strategies aimed at the improvement of plant phytoremediation potential using plant transformation and employing current achievements in nuclear and cytoplasmic genome transformation. Strategies for obtaining plants suitable for effective soil clean-up and tolerant to excessive concentrations of heavy metals are critically assessed. Promising directions in genetic manipulations, such as gene silencing and *cis*- and intragenesis, are also discussed. Moreover, the ways of overcoming disadvantages of phytoremediation using genetic transformation approach are proposed. The knowledge gathered here could be useful for designing new research aimed at biotechnological improvement of phytoremediation efficiency.

Keywords Environmental pollution · Genetic engineering · Genoremediation · Heavy metals · Hyperaccumulation · Transgenesis

Introduction

Dynamic development of industrial, urban, and agricultural aspects of human activities in the 20th century resulted in chemical pollution of the environment (Gavrilescu et al. 2015). In many parts of the world, the concentrations of numerous metallic elements in soil and water have increased to a level that endangers human health (Kelishadi et al. 2014). Release of small metal particles into the air also constitutes an environmental hazard of great concern (Serbula et al. 2012; Kularatne and de Oliveira; Freitas 2013). To prevent and treat hazardous pollution, wide range of techniques for heavy metal removal has been developed. However, the conventional physicochemical remediation is expensive

and detrimental to both microbiological life and soil properties (Fasenko and Edwards 2014; Singh and Santal 2015). Nowadays, apart from the traditional techniques involving mechanical or chemical treatment of a polluted substrate, science has supplied us with new opportunities of utilizing microorganisms (bioremediation), and plants (phytoremediation) to transform or remove toxic elements from the environment (Kang 2014; Mani and Kumar 2014; Gavrilescu et al. 2015). In numerous heavy metal-polluted locations, phytoremediation is a perspective technology of soil clean-up provided that plant genotypes with increased capability of stabilizing or extracting metals are employed. Phytoremediation techniques for treating metal-contaminated sites include phytoextraction, phytostabilization, rhizofiltration, and phytovolatilization (Ali et al. 2013; Lee 2013). Phytoextraction refers to an uptake of metals with further translocation and concentration in plant aboveground organs, whereas during phytostabilization, the toxic ions are maintained in the polluted substrate, being immobilized in the roots. Heavy metals are thus stabilized in the rhizosphere. In addition, the vegetation cover prevents dust blow and water erosion of the ground (Pilon-Smits and Pilon 2002; Singh and Santal 2015). During rhizofiltration, contaminants are the root

✉ Alina Wiszniewska
a.wiszniewska@ogr.ur.krakow.pl

¹ Institute of Plant Biology and Biotechnology, Faculty of Biotechnology and Horticulture, University of Agriculture in Krakow, Al. 29 Listopada 54, 31-425 Kraków, Poland

² Department of Botany, Faculty of Agriculture and Biology, Warsaw University of Life Sciences (SGGW), Nowoursynowska 159, Building 37, 02-776 Warsaw, Poland

absorbed, concentrated, or precipitated from contaminated water or polluted effluents by the root system of phytoremediating plant (Raskin et al. 1997). Phytoextraction and rhizofiltration require harvesting of the plant biomass with accumulated toxic metals. The plant material is then usually combusted, and the produced ash may be used for metal recycling (Jabeen et al. 2009; Ali et al. 2013). An innovative trend in using the harvested biomass from metal hyper-accumulating plants is a preparation of “eco-catalysts” for chemical industry (Escande et al. 2014, 2015). Plants are also capable of removing inorganic pollutants through volatilization. In this process, metals are biologically converted to gaseous forms and released into the atmosphere (Ruiz and Daniell 2009; Kang 2014).

The main progress in screening for efficient phytoremediating plants was achieved by selecting genotypes (ecotypes) growing naturally in contaminated sites (Reeves and Brooks 1983; Pollard et al. 2014; Muszyńska et al. 2015). Some progress was also achieved after a long-term selection of heavy metal-tolerant clones in *in vitro* and greenhouse conditions (Iori et al. 2012; Wiszniewska et al. 2015; Muszyńska and Hanus-Fajerska 2017). Further improvement of responsive genotypes by genetic engineering may help in making phytoremediation feasible for heavily heavy metal-polluted sites. Modifications may involve transformation of plants with different genes responsible for metal transport and homeostasis, defense response to oxidative stress, or xenobiotic detoxification. The latter genes are especially widespread in bacteria, and are a target of the so-called genomedi-ation, i.e., engineering of bacterial genes to increase their remediation potential, with subsequent incorporation into plant genome (Mani and Kumar 2014). Achievements of the last decades suggest that genetic engineering can be valuable tool in improving phytoremediation potential of the plant material belonging to different taxonomic groups. Our intention was to provide a comprehensive review on the current achievements in both nuclear and cytoplasmic genome transformation towards enhanced plant tolerance to heavy metals and their suitability for soil clean-up. In this review, we have discussed the effects of overexpression of genes encoding metal transporters, metal chelators, and components of antioxidative system. The main novelty of our paper is presentation of potential transformation-based solutions to overcome the most important phytoremediation drawbacks, limiting its broad exploitation. For the first time, we provided the comprehensive review of cytoplasmic genome transformation for improved plant potential to effective soil clean-up. In addition, we have presented and discussed genes that would enable to produce plant material useful not only for phytoextraction, but also for phytostabilization of contaminated areas. We discussed perspective directions in genetic manipulations aimed at enhancing phytoremediation of heavy metals, such as gene silencing and

cis- and intragenesis. The knowledge gathered here could be useful for designing novel research towards biotechnological improvement of phytoremediation efficiency.

Advantages and shortcomings of phytoremediation techniques

There are numerous advantages of phytoremediation technology. The most important are low costs in comparison with the traditional remediation technologies, eco-friendly and landscape-friendly nature, or significant reduction of the volume of contaminated material for disposal (Lee 2013; Wan et al. 2016). As a typical *in situ* technique, it may be applied to a broad range of sites, even those previously neglected due to high cost of conventional remediation technologies. However, there are also several limitations of this green technology. According to Ali et al. (2013), the main drawback is a long time required for a clean-up, what is related to limited capabilities of plants to accumulate high levels of heavy metals and to sustain normal growth in toxic environment. Plant remediation potential is also limited due to slow growth and low biomass accretion. Therefore, phytoremediation can be effective in sites with relatively low pollution level and is restricted to places where contaminants are reachable for the plant roots. In addition, phytoremediating plants are often unique ecotypes that inhabit specific ecosystems and their cultivation in other environmental conditions is difficult (Ali et al. 2013; Pollard et al. 2014). Finally, although public perception of phytoremediation is positive, improper biomass disposal may increase risk of food chain contamination with heavy metals. Modification of plant genotypes via genetic engineering could help to overcome main shortcomings of phytoremediation and make it more efficient (Pilon-Smits and Pilon 2002). Therefore, we attempted to suggest some potential transformation-based solutions to the most important phytoremediation limitations. In Table 1, we present main challenges that should be confronted during soil clean-up using phytotechnologies, and propose solutions based on genetic transformation, supported by examples of recent studies in this field.

Strategies for improved efficiency of phytoremediation

To improve phytoremediation traits using transgenesis, the most popular strategy is to introduce and overexpress the genes involved in the uptake, translocation, and sequestration of metals (Shukla et al. 2013; Mani and Kumar 2014; Das et al. 2016). Since metal metabolism in plant organism is extremely complex, there are numerous potential pathways that can be manipulated to increase metal accumulation in

Table 1 Phytoremediation challenges and selected examples of transformation-based solutions

Phytoremediation challenges (acc. to Ali et al. 2013)	Examples of transformation-based solutions	Effects	References
Reduction of time required for clean-up	Accelerating heavy metal uptake by plants overexpressing genes encoding novel metal transporters Transporters of phytochelator–metal complexes: yellow stripe (YS1) and yellow stripe-like (YLS) Co-transformation with multiple gene constructs	In <i>N. tabacum</i> , BJYSL7 gene from <i>B. juncea</i> enhanced accumulation of Cd and Ni In <i>A. thaliana</i> , SnYSL3 from <i>S. nigrum</i> gene is up-regulated in the excess of Cd and enhances translocation of Cd to the shoots In <i>A. thaliana</i> , dual-gene transformation with phytochelatin synthase gene from garlic (AsPCS1) and YCF1 (yeast vacuolar transporter) gene increased Cd and As accumulation properties	Wang et al. (2013) Feng et al. (2017) Guo et al. (2012)
Increased bioavailability of the contaminants in the soil	Modulation of heavy metal uptake via transformation with components of bacterial efflux system Transformation with genes encoding components of <i>Pseudomonas putida</i> efflux system: CzcA, CzcB and CzcC Increasing heavy metal uptake from the soil due to secretion of phytosiderophores Potential candidate genes: nicotianamine synthase (NAS) gene; nicotianamine amino transferase (NAAT) gene; deoxymygrinic acid synthase (DMAS) gene	In <i>N. tabacum</i> , increased bioavailability of Cd and gene variant-dependent differences in Cd accumulation in plant organs In hyperaccumulator <i>A. halterii</i> NAS regulates Zn bioavailability In <i>O. sativa</i> , HvNAS contributes to enhanced accumulation of Zn NAS and NAAT genes are expressed not only under Fe deficiency, but also in the presence of heavy metals (Cd, Pb and Ni)	Nesler et al. (2017) Tsednee et al. (2014) Masuda et al. (2008) Gupta and Singh (2017)
Accelerated growth rate and enhanced plant biomass accretion	Transformation-based organogenesis Transformation with rol genes from <i>Agrobacterium rhizogenes</i> Generation of hairy roots overexpressing modified copper resistance protein C (copC) gene from <i>Pseudomonas fluorescens</i> Stimulation of plant metabolism Transformation with heme oxygenase (HO) gene (regulation of phytochrome biosynthesis) Overexpression of isopenyl transferase (IPT) gene (increased synthesis of cytokinins)	In <i>S. americanus</i> , increased root biomass and plant tolerance to Pb and Cr In <i>N. tabacum</i> , increased root biomass, increased Cu accumulation and minimized toxicity symptoms In <i>B. napus</i> , higher plant biomass and increased tolerance of transformants to Hg stress, with decreased accumulation of Hg In <i>N. tabacum</i> , increased biomass accretion under Zn exposure	Alfaro-Saldana et al. (2016) Perez-Palacios et al. (2017) Shen et al. (2011) Pavlikova et al. (2014)
Increased plant tolerance to different growth condition (i.e., elevated level of contaminants, climate, and light intensity)	Transformation towards enhanced general stress defense mechanisms Genes encoding components of antioxidant system Genes encoding transcription factors related to various abiotic stresses (salinity, drought, and cold) Genes encoding elements of DNA repair system Genes encoding cytochromes	In <i>A. thaliana</i> , overexpression of OsGSTL2 (glutathione S-transferase) gene confers tolerance to various heavy metals and other abiotic stresses Gene encoding DRAB (drought-responsive element binding) transcription factor increased Cu tolerance in transgenic <i>N. tabacum</i> ; and Cd tolerance in transgenic <i>S. tuberosum</i> In <i>M. truncatula</i> , overexpression of MITdp2 α gene enhanced tolerance of plants exposed to Cu In <i>A. thaliana</i> , overexpression of CYP450-like gene (cytochrome P450) confers to tolerance to As, Cr, and Cd and other abiotic stresses	Kumar et al. (2013) Ban et al. (2011); Charfedine et al. (2017) Faè et al. (2014) Rai et al. (2015)

Table 1 (continued)

Phytoremediation challenges (acc. to Ali et al. 2013)	Examples of transformation-based solutions	Effects	References
Reduction of the risk of food chain contamination	The use of transgenic plants with increased ability to phytoremediation may itself accelerate soil clean-up and, as a result, the risk of food chain contamination can be diminished		

plant tissues, including metal mobilization and uptake by roots from the soil, formation of complexes with ligands and chelators, detoxification by deposition in vacuoles, and finally long-distance transport to shoots via symplast or apoplast (Nakamura et al. 2014; Das and Jayalekshmy 2015). The genes that are currently widely used to manipulate metal metabolism in plants are those that encode transporters of metal ions and metal-binding ligands (chelators).

Overexpression of genes encoding metal transporters

Plants developed versatile systems responsible for metal uptake and transportation, to supply cells with essential microelements, such as Cu^{2+} , Zn^{2+} , Fe^{2+} , and Co^{2+} (Williams et al. 2000). The plant genes encoding heavy metal transporters, usually represented by large gene families, are potential candidate genes for transformation towards improved phytoremediation potential. Manipulations involve enhancing of metal accumulation either in the roots for phytostabilization, or in the shoots for phytoextraction. Transformation may be also aimed at inhibiting heavy metal uptake and accumulation. Careful selection of suitable metal transporter genes is, therefore, linked with long-term purpose of the research, and some recent examples are briefly discussed in this section and presented in Table 2.

Tolerant transgenic plants may be obtained by manipulations concerning the efficiency of metal transport into a vacuole. There are several gene families that encode proteins functioning in cellular transport systems, and one of them is ATP-binding cassette (ABC) family, a large group of genes encoding proteins responsible for detoxification and ion-regulation processes (Martinoia et al. 2002). Protein transporters encoded by ABC genes were reported to be localized in the tonoplast of transgenic plant cells and to improve sequestration of metals in vacuolar lumen (Yazaki et al. 2006; Song et al. 2014). Advantageously, the effect of genetic transformation with ABC genes is directly linked with heavy metal mobility, and therefore, ABC genes could be overexpressed either to enhance translocation of mobile ions, like Cd and Cu, or to bind the non-mobile ones, like Pb, in the roots (Table 2). This way, plant ability to both phytoextraction and phytostabilization of heavy metals can be manipulated.

Genes belonging to cation diffusion facilitator (CDF) family encode various types of proteins called metal tolerance/transport protein (MTP), that enable cells to exclude excessive amounts of ions from cytoplasm. Plants overexpressing CDF genes produce elevated amounts of thiol compounds and, therefore, are able to efficiently sequester metal ions in the vacuoles by chelating them. This may result in hyperaccumulation of toxic ions in plant biomass. It was recently reported that during external Cd stress, overexpression of

Table 2 Recent examples of transformation using genes encoding metal transporters for increased efficiency of heavy metal phytoextraction and phyto-stabilization

Metal transporter family	Gene product	Gene origin and host	Effects	References
Phytoextraction				
ATP-binding cassette (ABC)	AtATM3	<i>A. thaliana</i> / <i>B. juncea</i>	Increased accumulation of Cd and Pb in the shoots	Bhuiyan et al. (2011)
ABC	YCF1	<i>S. cerevisiae</i> /Populus alba x P. tremula var. glandulosa	Increased accumulation of Cd and Zn in the shoots	Shim et al. (2013)
Iron regulated/ferroportin (IREG/FNP)	PgIREG1	Psychotria gabriellae/ <i>A. thaliana</i>	Increased accumulation of Ni in the shoots	Merlot et al. (2014)
Metal tolerance protein (MTP)	OsMTP1	<i>O. sativa</i> / <i>N. tabacum</i>	Increased accumulation of Cd in the shoots	Das et al. (2016)
Copper resistant protein (Cop)	copC	<i>Pseudomonas</i> sp./ <i>A. thaliana</i>	Hyperaccumulation of Cu in the shoots	Rodriguez-Lorente et al. (2012)
Phytostabilization				
P-1b-ATPase, CDF	AtHMA4	<i>A. thaliana</i> / <i>N. tabacum</i>	Restricted Cd transport from the roots to the shoots	Siemianowski et al. (2014)
ZIP	AtZIP1, AtMTP1	<i>A. thaliana</i> /Manihot esculenta	Zn accumulation in the roots, with restricted transport to the shoots	Gaitan-Solis et al. (2015)
ABC	YCF1	<i>S. cerevisiae</i> /Populus alba x P. tremula var. glandulosa	Increased accumulation of Pb in the roots	Shim et al. (2013)

OsMTP1 gene from indica rice (*Oryza sativa* L. cv. IR64) in tobacco stimulated hyperaccumulation of cadmium and enhanced tolerance and accumulation of arsenic upon exogenous. As stress, indicating broad substrate specificity of OsMTP1 (Das et al. 2016). It is an important finding, since this feature may be exploited in genetic modifications of plants towards enhanced co-tolerance to various trace metals in multi-contaminated sites.

Metal ion transporters are also products of large gene family encoding ZRT/IRT-related proteins (ZIP), which are engaged in the regulation of cytoplasmic transport of zinc and iron. In wild-type plants, these high affinity uptake systems are induced at transcriptional level by Zn or Fe deficiency (Eng et al. 1998). Majority of ZIP proteins form transmembrane ion channels with histidyl-rich metal-binding sites (Eng et al. 1998). Numerous recent studies focused on the exploitation of ZIP genes in crop biofortification with Zn and Fe (Tiong et al. 2015). However, these genes may be also overexpressed to improve phytoremediation of soil containing excessive amounts of heavy metals. In a model study, Conolly et al. (2002) revealed that overexpression of a gene encoding AtIRT1 in *A. thaliana* caused about 150% increase in accumulated amount of cadmium and zinc. More recent approach includes the use of metal transporters obtained from hyperaccumulator species, like *Nocca cearulescens*, which contributes to higher efficiency of soil clean-up. For instance, overexpression of NcZNT1 transgene encoding Zn transporter resulted in an enhanced accumulation of Zn and Cd in the transformed plants of *A. thaliana* (Lin et al. 2016). However, some ZIP proteins seem to be highly specific, and overexpression of their genes does not always result in improved accumulation of toxic heavy metals, even close analogues, like cadmium (Tiong et al. 2014). Therefore, another strategy exploited transformation with other genes encoding metal transporters and engaged in the uptake and transport of microelements. These genes may downregulate or upregulate the expression of ZIP genes, thus modifying the ability of transformed plants to accumulate toxic heavy metals. Usually, heavy metal uptake is significantly reduced due to increased concentration of microelements. This phenomenon can be useful in both phytoremediation of metal-contaminated soil, and crop biofortification in essential metals (Gong et al. 2015; Siemianowski et al. 2014).

Overexpression of genes encoding metal chelators

The main classes of metal chelators include phytochelatins (PCs), which are peptides enzymatically synthesized from glutathione, low molecular organic acids and amino acids (LMWOA), and metallothioneins (MT). Among these compounds, only metallothioneins are direct products of gene expression. In contrast, the level of phytochelatins and

organic and amino acids depends on the activity of enzymes involved in their biosynthesis, and thus genes encoding these enzymes should be manipulated.

Two key enzymes in the synthesis of PCs are phytochelatin synthase (PCS) and *c*-glutamyl cysteine synthetase (*c*-GCS) (Hirata et al. 2005). There are numerous examples of enhanced tolerance to heavy metals in transformed plants overexpressing phytochelatin synthase gene. Depending on the origin of the transgene, model plants of tobacco (*Nicotiana glauca* and *Nicotiana tabacum*) were either capable of greater accumulation of Cd and Pb, or were simply more tolerant to Cd, although without enhanced accumulation of this element (Huang et al. 2012; Chen et al. 2015). A serious concern of this approach is that various PCS differentially affect heavy metal transport within transformed plants. For example, PCS gene from *Populus tomentosa* reduced cadmium translocation to the aerial parts of tobacco (Chen et al. 2015), while PCS from aquatic macrophyte *Ceratophyllum demersum* enhanced this process (Shukla et al. 2013). An efficient way to overcome this issue could be co-transformation directed at multiple overexpression of genes involved in metal homeostasis. This approach may also increase the possibility to produce efficient hyperaccumulator. Guo et al. (2012) transformed *A. thaliana* with two genes: PCS, responsible for phytochelatin synthesis, and YCF1, ABC metal transporter, and obtained plants exhibiting co-tolerance to Cd and As, with increased bioaccumulation of these trace metals in vacuoles. Overexpression of ABC transporter gene reduced hypersensitivity reaction related to the accumulation of thiol-metal chelates, which, in turn, ameliorated plant tolerance. In addition, Zhao et al. (2014) proved that co-transformation with PCS and glutamyl cysteine synthetase (GCS) genes was beneficial to a transformation with PCS gene alone. They reported that overexpression of GCS enhanced the activity of PCS, resulting in higher production of phytochelatin. Due to this stimulatory effect, double transformants over-performed both wild-type plants and single transformants in relation to Cd accumulation and translocation.

Metal chelating properties were also reported for numerous organic low-molecular-weight compounds, such as organic acids and non-proteinogenic amino acids. Increased synthesis of these substances is often associated with plant tolerance to an excess of heavy metals (Lopez-Bucio et al. 2000). The genes encoding enzymes active in organic acid and amino acid synthesis can be also manipulated to improve phytoremediation process. A classical example could be an incorporation of HvNAS1 gene from *Hordeum vulgare* into *Arabidopsis*, reported by Kim et al. (2005). The gene encodes nicotianamine synthase, an enzyme that catalyzes trimerization of *S*-adenosyl methionine to nicotianamine, an amino acid capable of metal chelating. Due to increased biosynthesis of nicotianamine, the transformants

accumulated Ni more efficiently than wild-type plants and were capable of growing on Ni-rich serpentine soil without symptoms of Ni toxicity. This pioneer study revealed the role of nicotianamine in the mechanism of nickel detoxification by the formation of Ni–NA complexes that may be either deposited in vacuoles or translocated to the upper parts via phloem and xylem. Metals could be also chelated by organic acids, such as citric acid and oxalic acid. Although exudation of these compounds from the roots is specifically induced by aluminum, other stresses, including heavy metal exposure, also stimulate this process (Ding et al. 2014). In recent years, numerous genes related to organic acids exudation were identified, and, consequently, transformation experiments towards improved Al tolerance in plants were conducted (Wang et al. 2010, 2012; Zhou et al. 2014). Increased production of organic acids may be achieved by overexpression of the gene encoding enzymes catalyzing their metabolism, such as malate dehydrogenase, citrate synthase (CS), and phosphoenolpyruvate carboxylase (PEPC) (Wang et al. 2010, 2012). Another way may be the overexpression of genes encoding citrate transporters, such as those belonging to multidrug and toxic compound extrusion (MATE) family. In transformed plants, Al ions are effectively chelated in the root system, where citrate is released, and, therefore, toxic ions remain stabilized in the rhizosphere (Zhou et al. 2014). It was recently shown that overexpression of aluminum-induced protein (AIP) gene enhanced a tolerance of transformed *A. thaliana* plants to various heavy metals, including Cd and Cu (Jang et al. 2014). Based on these findings, a new strategy in phytoremediation area may be an exploitation of genes related to Al tolerance aimed at the enhancement of heavy metal chelation and stabilization.

Metallothioneins (MTs) are ubiquitous cysteine-rich proteins capable of high affinity coordination of heavy metal ions via cysteine residues gathered in Cys–X–Cys or Cys–Cys motifs. Plant metallothioneins are currently categorized into four types that slightly differ in a structure of cysteine-rich domains (Leszczyszyn et al. 2013). The role of plant MTs is generally attributed to the homeostasis of essential metals (Se, Zn, Ni, and Cu), and detoxification of xenobiotic metals (Cd, Hg, and Ag) (Tripathi et al. 2015). In recent years, great number of MT and MT-like gene sequences became available in databases and numerous studies reported on improved heavy metal tolerance in plants overexpressing MTs transgenes (Leszczyszyn et al. 2013). These reports were comprehensively reviewed and excellently discussed previously, for example by Eapen and Souza (2005), Kotrba et al. (2009), and Yadav et al. (2010). Therefore, here, we have just overviewed some very recent examples and presented them in Table 3. Instead, an interesting approach worth discussing is plant transformation with MTs genes isolated from halophytes.

Table 3 Recent examples of transformation using genes encoding metallothioneins (MTs) for improved heavy metal tolerance and accumulation in plants

Gene	Origin	Host	Effect	References
EhMT1	<i>Elsholtzia haichowensis</i>	<i>Nicotiana tabacum</i>	Higher tolerance to Cu and higher accumulation Cu in roots	Xia et al. (2012)
α MT1a	<i>Linum usitatissimum</i> L.	<i>Linum usitatissimum</i> L.	Overexpression of Cd-binding peptides enhanced accumulation of cadmium	Vrbova et al. (2015)
SpMT1	<i>Sedum plumbizincicola</i>	<i>Sedum plumbizincicola</i>	Elevated SpMTL transcript level might contribute to the trait of Cd hyperaccumulation and hyper-tolerance	Peng et al. (2017)
rgMT (type I)	<i>Oryza sativa</i> L.	<i>Sacharomyces cerevisiae</i> <i>A. thaliana</i>	Improved seed germination rates and increased biomass in the presence of heavy metal salts	Jin and Daniell (2014)
BcMT1 BcMT2	<i>Brassica campestris</i>	<i>Arabidopsis thaliana</i>	Enhanced tolerance to Cd and Cu and increased Cu concentration in the shoots; decreased Production of Cd- and Cu-induced ROS, thereby protecting plants from oxidative damage	Lu et al. (2015)
OsMT2c	<i>Oryza sativa</i> L.	<i>Arabidopsis thaliana</i>	Improved tolerance to Cu stress and increased ROS scavenging ability	Liu et al. (2015)
SaMT2	<i>Sedum alfredii</i>	<i>Saccharomyces cerevisiae</i> / <i>Nicotiana tabacum</i>	Increased Cd tolerance and accumulation	Zhang et al. (2014)
ThMT3	<i>Tamarix hispida</i>	<i>Salix matsudana</i>	Increased tolerance to Cu stress and higher NO productions (what contributed to the induction of adventitious roots under Cu stress)	Yang et al. (2015)
ThMT3	<i>Tamarix hispida</i>	<i>Saccharomyces cerevisiae</i>	Enhanced Cd, Zn, and Cu tolerance	Yang et al. (2011)

This group of plants includes species adapted to high salinity. Recent studies revealed that the mechanisms enabling plant growth under salt stress ameliorate also plant response to other abiotic stresses, including heavy metal treatment (Mariem et al. 2014; Taamalli et al. 2014; Guo et al. 2017). Metallothionein gene IIMt2a was successfully identified and characterized in a halophyte *Iris lactea* var. *chinensis* (Gu et al. 2014). Its subsequent incorporation into *Arabidopsis* genome resulted in higher tolerance of transgenic plants exposed to cadmium and copper (Gu et al. 2014, 2015). Heavy metal tolerance seemed to be associated with reduced production of reactive oxygen species (ROS), indicating high effectiveness of antioxidant defense system. Similar mechanisms led to increased heavy metal tolerance of tobacco transformed with SbMt2 gene from *Salicornia brachiata*, an extreme halophyte (Chaturvedi et al. 2014). Transgenic plants were capable of maintaining cellular homeostasis by modulating ROS scavenging/detoxification. Moreover, overexpression of SbMt2 gene was found to enhance Zn translocation into shoots, and therefore, it could be a good candidate gene for boosting stress tolerance and phytoremediation effectiveness via transgenesis (Chaturvedi et al. 2014). Future studies on genetic transformation towards improved phytoremediation potential could be significantly accelerated thanks to elucidating the mechanisms that allow extremophytes to survive in highly stressful conditions.

Strategies for improved heavy metal tolerance

Genetic transformation allows for obtaining plants exhibiting heavy metal resistance by manipulation with metabolic pathways responsible for stress defense reactions. At the cellular level, tolerance is usually manifested by an induction of mechanisms involved in DNA repair and radical scavenging. Using genetic manipulation of these two traits, one may further stimulate plant tolerance to toxic elements. The most common strategy in the studies on heavy metal stress is enhancement of plant antioxidant activity.

Overexpression of genes encoding components of antioxidant machinery

The oxidative damage is a consequence of various abiotic stresses, and occurs through the accumulation of elevated level of reactive oxygen species (ROS). Damaging effects of ROS to biomolecules, such as DNA, proteins, and lipids, can be minimized by antioxidant defense system including enzymatic mechanisms based mainly on superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), and glutathione *S*-transferase (GST) (Dixit et al. 2011; Hellou et al. 2012). It is, therefore, a popular approach in genetic transformation towards improved phytoremediation properties to increase the activity of antioxidant enzymes and ROS scavengers. Transformation experiments directed at

increasing general tolerance to abiotic stresses via overexpression of genes encoding antioxidant enzymes were conducted with various transgenes and host plants. However, the results related to the response to heavy metals were rather confusing. It was found that during an exposure to Cu, Cd, and As, transgenic plants overexpressing SOD, APX, and CAT genes generated lower amounts of ROS than wild-type plants, and the activity of antioxidant enzymes was highly elevated under stress (Lee et al. 2007; Guan et al. 2009; Gao et al. 2016). Nevertheless, the transgenic plants often exhibited both morphological and physiological disturbances such as growth retardation, abnormal root architecture, and reduction of photosynthetic efficiency at normal temperature condition (Iannone et al. 2015; Gao et al. 2016). Iannone et al. (2015) suggested that catalase did not play a crucial role in protecting tobacco plants against Cd toxicity. Instead, transgenic plants engineered for decreased activity of constitutive CAT (line CAT1AS) exposed to Cd were capable of activating alternative defense mechanisms against Cd stress, based mainly on increasing the activity of constitutive guaiacol peroxidase and ascorbate peroxidase. Moreover, Gao et al. (2016) reported that transformed plants of *Sedum alfredii* had reduced ability to accumulate cadmium ions than wild-type plants.

Contradictory results were also obtained for overexpression of glutathione *S*-transferase (GST) genes in plants subjected to heavy metal stress. These enzymes, possessing glutathione peroxidase activity, catalyze conjugation reactions between glutathione and a number of xenobiotics, and thereby contribute to the detoxification of free radicals (Hellou et al. 2012). Although transgenic tobacco plants expressing *Trichoderma virens* GST gene became more tolerant to cadmium ions and showed reduced lipid peroxidation in comparison with wild-type plants, Cd accumulation was not improved (Dixit et al. 2011). Growing transgenic alfalfa plants co-expressing GST and human cytochrome P450 2E1 genes turned out to be more efficient. The plants were more resistant to the toxic effects of heavy metals (Cd and Hg) and organic xenobiotic trichloroethylene and accumulated greater amounts of these pollutants than non-transgenic control plants (Zhang and Liu 2011; Zang et al. 2013). Such a combination of genes may be potentially used for phytoremediation of soils contaminated with a mixture of heavy metals and organic pollutants.

Attempts were also made to enhance synthesis of polyamines, since these compounds play an important role in the reduction of oxidative cell damage (Wen et al. 2010). For this purpose, a transgenic European pear (*Pyrus communis* L. 'Ballad') line was produced overexpressing apple spermidine synthase MdSPDS1 gene. After exposure to Cd and Zn, the transformants responded with enhanced synthesis and accumulation of polyamines as compared with wild-type plants. Beneficial effects were attributed to antioxidant activity of

spermidine and its ability to bind free metal ions (Wen et al. 2010). The essential role of polyamines in the reduction of stress symptoms in *Pyrus communis* was further confirmed during salt and cadmium treatment of transgenic lines containing MdSPDS1 in antisense orientation (Wen et al. 2011).

The main issue with enhancing heavy metal tolerance through overexpression of genes encoding antioxidant system is that there is not one defense mechanism against oxidative stress and the system of antioxidant response to the same heavy metal might differ in various plants and even in various organs and tissues within a plant organism. Schneider et al. (2013) provided an evidence for a cell type-specific antioxidant activity under Zn treatment and proved that epidermal cells exhibited higher accumulation of both glutathione and *S*-transferase proteins than the mesophyll tissue of *Noccaea caerulea* leaves. Therefore, the impact of tissue-specific gene expression should be taken into consideration during genetic modification of plants for biotechnological purposes. Besides, antioxidant enzymes may be identified as different isoforms of the same proteins, some of which may be up-regulated, while other down-regulated in response to heavy metals (Alvarez et al. 2009). Moreover, regulation of genes engaged in antioxidant reactions additionally depends on stress exposure, and thus, various antioxidant molecules are activated with the lapse of time (Ovečka and Takáč 2014).

The latest reports point at the possibility of stimulating plant heavy metal tolerance by overexpression of genes related to DNA repair and transcription. In this case, the mechanisms of tolerance are related to up-regulation of other genes, mainly those encoding antioxidant enzymes, such as cytosolic and chloroplastic isoforms of SOD and CAT, metallothioneins and metal transporters, as well as enzymes of DNA repair system (Faè et al. 2014; Charfeddine et al. 2017). Based on this, increased tolerance to Cu was obtained in transgenic *Medicago truncatula* overexpressing DNA repair gene MtTdp2 α , encoding tyrosyl-DNA phosphodiesterase 2 (Faè et al. 2014), and to Cu and Cd in potato overexpressing genes encoding transcription factors from drought-responsive element binding (DREB) family (Ban et al. 2011; Charfeddine et al. 2017). These examples showed that stress tolerance in plants is a very complex system and that efficient engineering of plants towards improved phytoremediation requires mechanisms involved in the cross-talk between heavy metals and plant response that remain to be explored in the future studies.

Transformation of chloroplast genome towards improved phytoremediation

Small chloroplast genome can be modified by genetic transformation, generating cells and plants with transgenic plastid genomes, which are referred to as transplastomic plants.

A specific feature of plastid transformation is homologous recombination that facilitates the site-specific alteration of endogenous plastid genes as well as the precisely targeted insertion of foreign genes into the plastid DNA (Bock 2015). This enables also overcoming common disadvantages of DNA transformation, such as gene silencing and position effect (Daniell et al. 2016). Homologous recombination has been used to analyze chloroplast gene functions and expression at all levels in vivo, and now, this method is nearly comparable to the techniques used for plant nuclear transformation (Lu et al. 2013; Bock 2015; Daniell et al. 2016). Benefits of chloroplast transformation include also the fact that codon optimization is not required to improve expression of bacterial transgenes, and therefore, any bacterial gene can be inserted in chloroplast genome (Quesada et al. 2005). An additional advantage of plastid transformation is minimizing the risk of transient gene transfer in the environments, since chloroplast DNA is maternally inherited and the dissemination of plastid DNA via pollen is rare phenomenon in plants (Maliga 2002). Chloroplast transformation is especially suitable for the development of transgenic plants when multiple genes are required for effective phytoremediation, and therefore, it is an ideal approach for mercury phytoremediation. Pioneer studies revealed that transformation of tobacco chloroplast genome by native bacterial genes *merA* and *merB* induced plant resistance to very high concentrations, up to 400 mM, of phenylmercuric acetate (PMA). Transgenic plants not only survived such high concentrations of PMA but also grew better than control untreated plants (Ruiz et al. 2003; Hussein et al. 2007). Although it was reported that organic Hg is transported in plants more efficiently than inorganic forms of mercury (Kumar et al. 2004), transgenic tobacco volatilized Hg⁰ efficiently, independently of the form of Hg in the soil, confirming that both mercuric ion reductase and organomercuriallyase were active in transgenic chloroplasts (Hussein et al. 2007). Nevertheless, a disadvantage of the described strategy is that elemental mercury is volatilized from the cell and released back into the environment, where it can be converted into highly toxic forms. For this reason, attempts had been made to stimulate mercury accumulation in plant tissues for their subsequent utilization (Ruiz et al. 2011; Zhang et al. 2013). Ruiz et al. (2011) reported on a development of a transplastomic system to express murine metallothionein gene (MT1) that enabled accumulation of mercury in high concentrations within tobacco cells. The transgenic lines were resistant up to 20 µM mercury, maintained high chlorophyll content and biomass, and actively translocated Hg to the leaves, indicating increased phytoremediation capacity. Chloroplast transformation may also increase plant tolerance to stress induced by excessive concentration of copper. Jin and Daniell (2014) reported that overexpression of γ -tocopherol methyltransferase gene (γ -TMT) in transformed

tobacco chloroplasts resulted in accumulation of the enzyme in the inner envelope membrane. Due to increased activity of the enzyme, a concentration of α -tocopherol in transplastomic seeds was twice as high as in wild-type ones. As a consequence, transplastomic plants grew better in the presence of excessive amounts of copper, exhibiting increased tolerance related to antioxidant defense (Jin and Daniell 2014). Martret et al. (2011) also attempted to stimulate plant antioxidant system under cadmium and zinc treatment by transformation of tobacco chloroplast genome with three genes encoding antioxidant enzymes: dehydroascorbate reductase (DHAR), glutathione *S*-transferase (GST), and glutathione reductase (GR). Although transplastomic plants were less sensitive to various stress conditions, such as increased salinity and chilling, chloroplast transformation did not ameliorate plants response to heavy metal exposure. This illustrates the fact that in some cases, nuclear transformation has more pronounced impact on antioxidant system than chloroplast transformation. Therefore, screening for appropriate genes is essential to successfully transform plastids towards heavy metal tolerance. A good candidate gene is BrMT1 encoding *Brassica rapa* type-1 metallothionein. This gene was successfully targeted to the chloroplasts of *Arabidopsis* and its overexpression in either the chloroplasts or the cytosol allowed for effective detoxification of cadmium in transplastomic plants (Kim et al. 2007). In particular, the chloroplast-targeted BrMT1 was associated with a significant reduction in chlorosis and accumulation of H₂O₂. Considering efficiency of plastid transformation, there are some concerns about the functioning of transgenic chloroplasts in the roots, where are naturally absent. Since root system plays a crucial role in phytoremediation, Davison (2005) doubts whether chloroplast transformation would be effective tool for enhanced heavy metal uptake by this organ. This problem could be by-passed by elaboration of transformation protocols for other types of plastids, especially leucoplasts, that are present also in organs that are not active in photosynthesis. Recently, similar approach has been successfully applied in creating transplastomic *Nicotiana* plants exhibiting elevated tolerance to multiple biotic and abiotic stresses (Chen et al. 2014).

Miscellaneous directions of transformation towards improved phytoremediation

Algal species, especially microalga, have great ability to absorb and transform pollutants, including heavy metals. These organisms synthesize both phytochelatin and metallothioneins, and, therefore, are capable of chelating and efficiently detoxifying toxic elements (Perales-Vela et al. 2006). However, the algae have not been widely studied and exploited for soil remediation purposes, since they naturally

dwelling in aquatic environments. Nevertheless, due to their unicellular structure, algal cells may serve as a tool in fundamental research studying cellular mechanisms of heavy metal tolerance. They can also be a source of novel genes to be used in plant transformation. A good example is *Chlamydomonas*, in the genome of which at least 11 unique gene families known to encode metal transporters were described (Rubinelli et al. 2002). Another interesting feature is specific cell wall structure in alga, which contains sulphated oligosaccharides playing a role in heavy metal binding (Xue et al. 1988). Investigation in this field opens new possibilities of engineering plant cell wall proteins to increase their metal-binding capacity.

An emerging transformation approach for increased heavy metal resistance may become gene silencing. Biological basis for this process is RNA interference, a conserved defense mechanism, in which small RNA molecules suppress gene expression by interfering with the translation of target mRNA (Saurabh et al. 2014). This technique was applied to improve crop quality in several important traits, such as allergen elimination, altered plant phenotype, biofortification, seedless fruit production, and more (see review of Saurabh et al. 2014). Gene silencing allows for avoiding heavy metal accumulation, as reported by Li et al. (2007), where Cd content in grains was drastically reduced by silencing phytochelatin synthase gene. However, gene silencing via RNA interference may also facilitate heavy metal uptake, resulting in hyperaccumulation of toxic elements. Dhanker et al. (2006) reported that *Arabidopsis* plants silenced for arsenate reductase gene were capable of efficiently translocating arsenic to the shoots, in contrast with wild-type plants (Dhankher et al. 2006). Enhanced Cd translocation to the shoots was also achieved in rice by silencing the gene encoding root-localized Cd-transporter OsNRAMP5 (Takahashi et al. 2014). Such an approach would help to intensify phytoextraction of metallic pollutants from the soil. As gene silencing technology has numerous benefits, such as precision, efficiency, stability, and flexibility (Saurabh et al. 2014), substantial increase in phytoremediation efficiency can be expected in the nearest future.

Future prospects to overcome transformation concerns

Genetic engineering is a powerful tool that may accelerate development of new plant lines with traits desirable in phytoremediation. However, there have been public concerns about testing transgenic lines in field conditions, and therefore, no genetically modified plants have been tested in large-scale field experiments for their specific potential in heavy metal remediation. Thus, also the potential risks associated with outcrossing with any wild relatives

or uncontrolled spread due to higher tolerance level have never been clearly evaluated. The risk of gene escape with engineered hyperaccumulator lines can be reduced if plant material is applied in isolated industrial districts instead of agricultural areas. A novel solution can be an application of cisgenesis and intragenesis in genetic engineering of phytoremediating plants. These new tools in plant genetic modification were successfully applied in improving crop quality and enhancing tolerance to biotic stresses (Espinoza et al. 2013). In both methods, plants are modified with the copies of natural genes: in cisgenesis, the construct must contain complete genetic elements from a sexually compatible gene donor (promoter, coding sequence including introns, and terminator), while in intragenesis, the constructs may contain genetic elements from different genes within a sexually compatible gene pool (Espinoza et al. 2013). Cisgenic and intragenic approach may be especially valuable in breeding programs that would facilitate biomass production and growth of natural hyperaccumulators. Plants belonging to Brassicaceae family, such as *Arabidopsis* or *Thlaspi*, are good candidates for genetic modifications due to scientific knowledge on their genetics and their close phylogenetic relationships. Genetic foundations of heavy metal tolerance were extensively studied in this family and several ecotypes of hyperaccumulator species were identified (Marmiroli et al. 2006). Recent findings in genetic mapping in maize may facilitate a production of cisgenic plants with enhanced/limited ability to accumulate cadmium in leaf tissues, which can be exploited in phytoremediation of contaminated soil (Zdunić et al. 2014). Although, to date, there have been no reports exploiting cisgenesis and/or intragenesis in modulating plant responses to heavy metal stress, these novel techniques provide an excellent opportunity to overcome concerns with transgenesis and accelerate studies on genetic manipulations towards increased efficiency of phytoremediation.

Conclusions

In this review, we have comprehensively discussed recent achievements and novel directions of plant transformation towards enhanced heavy metal tolerance and phytoremediation. We also proposed some potential transformation-based solutions to the most important phytoremediation drawbacks. In this view, plant transformation may facilitate soil clean-up for sustainable environmental protection. Since genetic engineering may be an extremely powerful tool in environmental studies, especially taking into account that new technologies, such as genome editing and genome engineering, are still emerging, the knowledge gathered here could be useful in designing new research towards biotechnological improvement of phytoremediation efficiency.

Acknowledgements This research was financed by the Ministry of Science and Higher Education of the Republic of Poland to the Institute of Plant Biology and Biotechnology, University of Agriculture in Kraków (Grant no. DS 3500).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alfaro-Saldaña EF, Pérez-Molpe-Balch E, Santos-Díaz MS (2016) Generation of transformed roots of *Scirpus americanus Pers.* and study of their potential to remove Pb²⁺ and Cr³⁺. *Plant Cell Tissue Organ Cult* 127(1):15–24
- Ali H, Khan E, Sajad MA (2013) Phytoremediation of heavy metals—concepts and applications. *Chemosphere* 91:869–881
- Alvarez S, Berla MB, Sheffield J, Cahoon RE, Jez JM, Hicks LM (2009) Comprehensive analysis of the *Brassica juncea* root proteome in response to cadmium exposure by complementary proteomic approaches. *Proteomics* 9:2419–2431
- Ban Q, Liu G, Wang Y (2011) A DREB gene from *Limonium bicolor* mediates molecular and physiological responses to copper stress in transgenic tobacco. *J Plant Physiol* 168:449–458
- Bhuiyan MSU, Min SR, Jeong WJ, Sultana S, Choi KS, Lee Y, Liu JR (2011) Overexpression of AtATM3 in *Brassica juncea* confers enhanced heavy metal tolerance and accumulation. *Plant Cell Tissue Organ Cult* 107(1):69–77
- Bock R (2015) Engineering plastid genomes: methods, tools, and applications in basic research and biotechnology. *Ann Rev Plant Biol* 66:211–241
- Charfeddine M, Charfeddine S, Bouaziz D, Messaoud RB, Bouzid RG (2017) The effect of cadmium on transgenic potato (*Solanum tuberosum*) plants overexpressing the StDREB transcription factors. *Plant Cell Tissue Organ Cult* 128(3):521–541
- Chaturvedi AK, Patel MK, Mishra A, Tiwari V, Jha B (2014) The SbMT-2 gene from a halophyte confers abiotic stress tolerance and modulates ROS scavenging in transgenic tobacco. *PLoS One* 9(10):e111379
- Chen PJ, Senthilkumar R, Jane WN, He Y, Tian Z, Yeh KW (2014) Transplastomic *Nicotiana benthamiana* plants expressing multiple defence genes encoding protease inhibitors and chitinase display broad-spectrum resistance against insects, pathogens and abiotic stresses. *Plant Biotechnol J* 12(4):503–515
- Chen Y, Liu Y, Ding Y, Wang X, Xu J (2015) Overexpression of PtPCS enhances cadmium tolerance and cadmium accumulation in tobacco. *Plant Cell Tissue Organ Cult* 121(2):389–396
- Conolly EL, Fett JP, Guerinot ML (2002) Expression of the IRT1 metal transporter is controlled by metals at the levels of transcript and protein accumulation. *Plant Cell* 14(6):1347–1357
- Daniell H, Lin C, Yu M, Chang W (2016) Chloroplast genomes: diversity, evolution and applications in genetic engineering. *Genome Biol* 17:134
- Das R, Jayalekshmy VG (2015) Mechanism of heavy metal tolerance and improvement of tolerance in crop plants. *J Glob Biosci* 4(7):2678–2698
- Das N, Bhattacharya S, Maiti MK (2016) Enhanced cadmium accumulation and tolerance in transgenic tobacco overexpressing rice metal tolerance protein gene OsMTP1 is promising for phytoremediation. *Plant Physiol Biochem* 105:297–309
- Davison J (2005) Risk mitigation of genetically modified bacteria and plants designed for bioremediation. *J Ind Microbiol Biotechnol* 32(11–12):639–650
- Dhankher OP, Rosen BP, McKinney EC, Meagher RB (2006) Hyperaccumulation of arsenic in the shoots of *Arabidopsis* silenced for arsenate reductase (ACR2). *Proc Natl Acad Sci USA* 103(14):5413–5418
- Ding H, Wen D, Fu Z, Qian H (2014) The secretion of organic acids is also regulated by factors other than aluminum. *Environ Monit Assess* 186(2):1123–1131
- Dixit P, Mukherjee PK, Ramachandran V, Eapen S (2011) Glutathione transferase from *Trichoderma virens* enhances cadmium tolerance without enhancing its accumulation in transgenic *Nicotiana tabacum*. *PLoS One* 6(1):e16360. <https://doi.org/10.1371/journal.pone.0016360>
- Eapen S, D'Souza SF (2005) Prospect of genetic engineering of plants for phytoremediation of toxic metals. *Biotechnol Adv* 23:97–114
- Eng BH, Guerinot ML, Eide D, Saier Jr MH (1998) Sequence analyses and phylogenetic characterization of the ZIP family of metal ion transport proteins. *J Membr Biol* 166(1):1–7
- Escande V, Olszewski TK, Grison C (2014) Preparation of ecological catalysts derived from Zn hyperaccumulation plants and their catalytic activity in Diels–Alder reaction. *CR Chim* 17:731–737
- Escande V, Olszewski T, Grison C (2015) From biodiversity to catalytic diversity: how to control the reaction mechanism by the nature of metallophytes. *Environ Sci Pollut Res* 22:5653–5666
- Espinoza C, Schlechter R, Herrera D, Torres E, Serrano A, Medina C, Arce-Johnson P (2013) Cisgenesis and intragenesis: new tools for improving crops. *Biol Res* 46(4):323–331
- Faè M, Balestrazzi A, Confalonieri M, Donà M, Macovei A, Valassi A, Carbonera D (2014) Copper-mediated genotoxic stress is attenuated by the overexpression of the DNA repair gene MtTdp2α (tyrosyl-DNA phosphodiesterase 2) in *Medicago truncatula* plants. *Plant Cell Rep* 33(7):1071–1080
- Fasenko E, Edwards R (2014) Plant synthetic biology: a new platform for industrial biotechnology. *J Exp Bot* 65(8):1927–1937
- Feng S, Tan J, Zhang Y, Liang S, Xiang S, Wang H, Chai T (2017) Isolation and characterization of a novel cadmium-regulated Yellow Stripe-Like transporter (SnYSL3) in *Solanum nigrum*. *Plant Cell Rep* 36:281–296
- Gaitán-Solís E, Taylor NJ, Siritunga D, Stevens W, Schachtman DP (2015) Overexpression of the transporters AtZIP1 and AtMTP1 in cassava changes zinc accumulation and partitioning. *Front Plant Sci* 6:492
- Gao X, Ai WL, Gong H, Cui LJ, Chen BX, Luo HY, Qiu BS (2016) Transgenic NtFeSOD *Sedum alfredii* plants exhibited profound growth impairments and better relative tolerance to long-term abiotic stresses. *Plant Biotechnol Rep* 10(2):117–128
- Gavrilescu M, Demnerova K, Aamand J, Agathos S, Fava F (2015) Emerging pollutants in the environment: present and future challenges in biomonitoring, ecological risks and bioremediation. *New Biotechnol* 32:147–156
- Gong X, Yin L, Chen J, Gu C (2015) Overexpression of the iron transporter NtPIC1 in tobacco mediates tolerance to cadmium. *Plant Cell Rep* 34(11):1963–1973
- Gu CS, Liu LQ, Zhao YH, Deng YM, Zhu XD, Huang SZ (2014) Overexpression of *Iris lactea var. chinensis* metallothionein IIMT2a enhances cadmium tolerance in *Arabidopsis thaliana*. *Ecotoxicol Environ Saf* 105:22–28
- Gu CS, Liu LQ, Deng YM, Zhu XD, Huang SZ, Lu XQ (2015) The heterologous expression of the *Iris lactea var. chinensis* type 2 metallothionein IIMT2b gene enhances copper tolerance in *Arabidopsis thaliana*. *Bull Environ Contam Toxicol* 94(2):247–253

- Guan Z, Chai T, Zhang Y, Xu J, Wei W (2009) Enhancement of Cd tolerance in transgenic tobacco plants overexpressing a Cd-induced catalase cDNA. *Chemosphere* 76(5):623–630
- Guo J, Xu W, Ma M (2012) The assembly of metals chelation by thiols and vacuolar compartmentalization conferred increased tolerance and accumulation of cadmium and arsenic in transgenic *Arabidopsis thaliana*. *J Hazard Mater* 199–200:309–313
- Guo Q, Meng L, Zhang YN, Mao PC, Tian XX, Li SS, Zhang L (2017) Antioxidative systems, metal ion homeostasis and cadmium distribution in *Iris lactea* exposed to cadmium stress. *Ecotoxicol Environ Saf* 139:50–55
- Gupta CK, Singh B (2017) Uninhibited biosynthesis and release of phytosiderophores in the presence of heavy metal (HM) favors HM remediation. *Environ Sci Pollut Res* 24(10):9407–9416
- Hellou J, Ross NW, Moon TW (2012) Glutathione, glutathione S-transferase and glutathione conjugates, complementary markers of oxidative stress in aquatic biota. *Environ Sci Pollut Res* 19:2007–2023
- Hirata K, Tsuji N, Miyamoto K (2005) Biosynthetic regulation of phytochelatin, heavy metal-binding peptides. *J Biosci Bioeng* 100:593–599
- Huang J, Zhang Y, Peng J-S, Zhong C, Yi H-Y, Ow DW, Gong Y-M (2012) Fission yeast HMT1 lowers seed cadmium through phytochelatin-dependent vacuolar sequestration in *Arabidopsis*. *Plant Physiol* 158:1779–1788
- Hussein HS, Ruiz ON, Terry N, Daniell H (2007) Phytoremediation of mercury and organomercurials in chloroplast transgenic plants: enhanced root uptake, translocation to shoot and volatilization. *Environ Sci Technol* 41:8439–8446
- Iannone MF, Groppa MD, Benavides MP (2015) Cadmium induces different biochemical responses in wild type and catalase-deficient tobacco plants. *Environ Exp Bot* 109:201–211
- Iori V, Pietrini F, Massacci A, Zacchini M (2012) Induction of metal binding compounds and antioxidative defense in callus cultures of two black poplar (*P. nigra*) clones with different tolerance to cadmium. *Plant Cell Tissue Organ Cult* 108(1):17–26
- Jabeen R, Ahmad A, Iqbal M (2009) Phytoremediation of heavy metals: physiological and molecular mechanisms. *Bot Rev* 75:339–364
- Jang MG, Kim YJ, Jang GH, Sukweenadhi J, Kwon WS, Yang DC (2014) Ectopic overexpression of the aluminum-induced protein gene from *Panax ginseng* enhances heavy metal tolerance in transgenic *Arabidopsis*. *Plant Cell Tissue Organ Cult* 119(1):95–106
- Jin S, Daniell H (2014) Expression of gamma-tocopherol methyltransferase in chloroplasts results in massive proliferation of the inner envelope membrane and decreases susceptibility to salt and metal-induced oxidative stresses by reducing reactive oxygen species. *Plant Biotechnol J* 12:1274–1285
- Kang JW (2014) Removing environmental organic pollutants with bioremediation and phytoremediation. *Biotechnol Lett* 36(6):1129–1139
- Kelishadi R, Moeini R, Poursafa P, Farajian S, Yousefy H, Okhovat-Souraki A (2014) Independent association between air pollutants and vitamin D deficiency in young children in Isfahan, Iran. *Paediatr Int Child Health* 34:50–55
- Kim S, Takahashi M, Higuchi K, Tsunoda K, Nakanishi H, Yoshimura E, Nishizawa NK (2005) Increased nicotianamine biosynthesis confers enhanced tolerance of high levels of metals, in particular nickel, to plants. *Plant Cell Physiol* 46(11):1809–1818
- Kim SH, Lee HS, Song WY, Choi KS, Hur Y (2007) Chloroplast targeted BrMT1 (*Brassica rapa* Type-I Metallothionein) enhances resistance to cadmium and ROS in transgenic *Arabidopsis* plants. *J Plant Biol* 50(1):1–7
- Kotrba P, Najmanowa J, Macek T, Ruml T, Mackova M (2009) Genetically modified plants in phytoremediation of heavy metal and metalloids soil and sediment pollution. *Biotechnol Adv* 29:799–810
- Kumar S, Asif MH, Chakrabarty D, Tripathi RD, Dubey RS, Trivedi PK (2013) Expression of a rice Lambda class of glutathione S-transferase, OsGSTL2, in arabidopsis provides tolerance to heavy metal and other abiotic stresses. *J Hazard Mater* 248:228–237
- Kularatne KIA, De Freitas CR (2013) Epiphytic lichens as biomonitors of airborne heavy metal pollution. *Environ Exp Bot* 88:24–32
- Kumar S, Dhingra A, Daniell H (2004) Plastid-expressed betaine aldehyde dehydrogenase gene in carrot cultured cells, roots, and leaves confers enhanced salt tolerance. *Plant Physiol* 136:2843–2854
- Lee SH, Ahsan N, Lee KW, Kim DH, Lee DG, Kwak SS, Lee BH (2007) Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses. *J Plant Physiol* 164(12):1626–1638
- Lee JH (2013) An overview of phytoremediation as a potentially promising technology for environmental pollution control. *Biotechnol Bioprocess Eng* 18(3): 431–439
- Leszczyszyn OI, Imam HT, Blindauer CA (2013) Diversity and distribution of plant metallothioneins: a review of structure, properties and functions. *Metallomics* 5(9):1146–1169
- Li JC, Guo JB, Xu WZ, Ma M (2007) RNA interference mediated silencing of phytochelatin synthase gene reduce cadmium accumulation in rice seeds. *J Integr Plant Biol* 49(7):1032–1037
- Lin Y-F, Hassan Z, Talukdar S, Schat H, Aarts MGM (2016) Expression of the ZNT1 zinc transporter from the metal hyperaccumulator *Noccaea caerulea* confers enhanced zinc and cadmium tolerance and accumulation to *Arabidopsis thaliana*. *PLoS One* 11(3):e0149750. <https://doi.org/10.1371/journal.pone.0149750>
- Liu J, Shi X, Qiana M, Zhenga Lm Lianb C, Yan Xiaa Y, Shen Z (2015) Copper-induced hydrogen peroxide upregulation of a metallothionein gene, OsMT2c, from *Oryza sativa* L.confers copper tolerance in *Arabidopsis thaliana*. *J Hazard Mater* 294:99–108
- Lopez-Bucio J, Nieto-Jacobo MF, Ramirez-Rodriguez V, Herrera-Estrella L (2000) Organic acid metabolism in plants: from adaptive physiology to transgenic varieties for cultivation in extreme soils. *Plant Sci* 160(1):1–13
- Lu Y, Rijzaani H, Karcher D, Ruf S, Bock R (2013) Efficient metabolic pathway engineering in transgenic tobacco and tomato plastids with synthetic multigene operons. *Proc Natl Acad Sci USA* 110:E623–E632
- Lu Y, Deng X, Quan L, Xia Y, Shen Z (2015) Metallothioneins BcMT1 and BcMT2 from *Brassica campestris* enhance tolerance to cadmium and copper and decrease production of reactive oxygen species in *Arabidopsis thaliana*. *Plant Soil* 367:507–519
- Maliga P (2002) Engineering the plastid genome of higher plants. *Curr Opin Plant Biol* 5(2):164–172
- Mani D, Kumar C (2014) Biotechnological advances in bioremediation of heavy metals contaminated ecosystems: an overview with special reference to phytoremediation. *Int J Environ Technol* 11:843–872
- Mariem W, Kilani BR, Benet G, Abdelbasset L, Stanley L, Charlotte P, Tahar G (2014) How does NaCl improve tolerance to cadmium in the halophyte *Sesuvium portulacastrum*? *Chemosphere* 117:243–250
- Marmiroli N, Marmiroli M, Maestri E (2006) Phytoremediation and phytotechnologies: a review for the present and the future. In: Soil and water pollution monitoring, protection and remediation. Springer, Dordrecht, pp 403–416
- Martinoia E, Klein M, Geisler M, Bovet L, Forestier C, Kolukisaoglu Ü, Schulz B (2002) Multifunctionality of plant ABC transporters—more than just detoxifiers. *Planta* 214(3):345–355

- Martret BL, Poage M, Shiel K, Nugent GD, Dix PJ (2011) Tobacco chloroplast transformants expressing genes encoding dehydroascorbate reductase, glutathione reductase, and glutathione-S-transferase, exhibit altered anti-oxidant metabolism and improved abiotic stress tolerance. *Plant Biotechnol J* 9:661–673
- Masuda H, Suzuki M, Morikawa KC, Kobayashi T, Nakanishi H, Takahashi M, Saigusa M, Mori S, Nishizawa NK (2008) Increase in iron and zinc concentrations in rice grains via the introduction of barley genes involved in phytosiderophore synthesis. *Rice* 1:100–108
- Merlot S, Hannibal L, Martins S, Martinelli L, Amir H, Lebrun M, Thomine S (2014) The metal transporter PgIREG1 from the hyperaccumulator *Psychotriagabriellae* is a candidate gene for nickel tolerance and accumulation. *J Exp Bot* 65(6):1551–1564
- Muszyńska E, Hanus-Fajerska E (2017) In vitro multiplication of *Dianthus carthusianorum* calamine ecotype with the aim to revegetate and stabilize polluted wastes. *Plant Cell Tissue Organ Cult* 128:631–640
- Muszyńska E, Hanus-Fajerska E, Ciarkowska K (2015) Studies on *Gypsophila fastigiata* parameters verifying its suitability to reclamation of post-flotation Zn-Pb wastes. *Geol Geophys Environ* 41(1):17–24
- Nakamura M, Ochiai T, Noji M, Ogura Y, Suzuki K, Yoshimoto N, Yamazaki M, Saito K (2014) An improved tolerance to cadmium by overexpression of two genes for cysteine synthesis in tobacco. *Plant Biotechnol* 31:141–147
- Nesler A, DalCorso G, Fasani E, Manara A, Di Sansebastiano GP, Argese E, Furini A (2017) Functional components of the bacterial CzcCBA efflux system reduce cadmium uptake and accumulation in transgenic tobacco plants. *New Biotechnol* 35:54–61
- Ovečka M, Takáč T (2014) Managing heavy metal toxicity stress in plants: biological and biotechnological tools. *Biotechnol Adv* 32:73–86
- Pavlíková D, Pavlík M, Procházková D, Zemanová V, Hnilička F, Wilhelmová N (2014) Nitrogen metabolism and gas exchange parameters associated with zinc stress in tobacco expressing an IPT gene for cytokinin synthesis. *J Plant Physiol* 171(7):559–564
- Peng JS, Ding G, Meng S, Yi HY, Gong JM (2017) Enhanced metal tolerance correlates with heterotypic variation in SpMTL, a metallothionein-like protein from the hyperaccumulator *Sedum plumbizincicola*. *Plant Cell Environ*. <https://doi.org/10.1111/pce.12929>
- Perales-Vela HV, Pena-Castro JM, Canizares-Villanueva RO (2006) Heavy metal detoxification in eukaryotic microalgae. *Chemosphere* 64(1):1–10
- Pérez-Palacios P, Agostini E, Ibáñez SG, Talano MA, Rodríguez-Llorente ID, Caviedes MA, Pajuelo E (2017) Removal of copper from aqueous solutions by rhizofiltration using genetically modified hairy roots expressing a bacterial Cu-binding protein. *Environ Technol* 38(22):2877–2888
- Pilon-Smits E, Pilon M (2002) Phytoremediation of metal using transgenic plants. *Crit Rev Plant Sci* 21(5):439–456
- Pollard AJ, Reeves RD, Baker AJ (2014) Facultative hyperaccumulation of heavy metals and metalloids. *Plant Sci* 217:8–17
- Quesada T, Ruiz ON, Daniell H (2005) Characterization of heterologous multigene operons in transgenic chloroplasts: transcription, processing and translation. *Plant Physiol* 138:1–17
- Rai A, Singh R, Shirke PA, Tripathi RD, Trivedi PK, Chakrabarty D (2015) Expression of rice CYP450-like gene (Os08g01480) in *Arabidopsis* modulates regulatory network leading to heavy metal and other abiotic stress tolerance. *PLoS One* 10(9):e0138574
- Raskin I, Smith RD, Salt DE (1997) Phytoremediation of metals: using plants to remove pollutants from the environment. *Curr Opin Biotechnol* 8(2):221–226
- Reeves RD, Brooks RR (1983) Hyperaccumulation of lead and zinc by two metallophytes from mining areas of Central Europe. *Environ Pollut A* 31(4):277–285
- Rodríguez-Llorente ID, Lafuente A, Doukkali B, Caviedes MA, Pajuelo E (2012) Engineering copper hyperaccumulation in plants by expressing a prokaryotic cop C gene. *Environ Sci Technol* 46(21):12088–12097
- Rubinelli P, Siripornadulsil S, Gao-Rubinelli F, Sayre RT (2002) Cadmium- and iron-stress-inducible gene expression in the green alga *Chlamydomonas reinhardtii*: evidence for H43 protein function in iron assimilation. *Planta* 215(1):1–13
- Ruiz ON, Daniell H (2009) Genetic engineering to enhance mercury phytoremediation. *Curr Opin Biotechnol* 20:213–219
- Ruiz ON, Hussein HS, Terry N, Daniell H (2003) Phytoremediation of organomercurial compounds via chloroplast genetic engineering. *Plant Physiol* 132:1344–1352
- Ruiz ON, Alvarez D, Torres C, Roman L, Daniell H (2011) Metallothionein expression in chloroplasts enhances mercury accumulation and phytoremediation capability. *Plant Biotechnol J* 9:609–617
- Saurabh S, Vidyarthi AS, Prasad D (2014) RNA interference: concept to reality in crop improvement. *Planta* 239(3):543–564
- Schneider T, Persson DP, Husted S, Schellenberg M, Gehrig P, Lee Y, Martinoia E, Schjoerring JK, Meyer S (2013) A proteomics approach to investigate the process of Zn hyperaccumulation in *Noccaea caerulea* (J & Presl C) F.K. Meyer. *Plant J* 73:131–142
- Serbula SM, Miljkovic DD, Kovacevic RM, Ilic AA (2012) Assessment of airborne heavy metal pollution using plant parts and topsoil. *Ecotoxicol Environ Saf* 76:209–214
- Shen Q, Jiang M, Li H, Che LL, Yang ZM (2011) Expression of a *Brassica napus* hemoxygenase confers plant tolerance to mercury toxicity. *Plant Cell Environ* 34(5):752–763
- Shim D, Kim S, Choi YI, Song WY, Park J, Youk ES, Jeong S-C, Martinoia E, Noh E-W, Lee Y (2013) Transgenic poplar trees expressing yeast cadmium factor 1 exhibit the characteristics necessary for the phytoremediation of mine tailing soil. *Chemosphere* 90(4):1478–1486
- Shukla D, Kesari R, Tiwari M, Dwivedi S, Tripathi RD, Nath P, Trivedi PK (2013) Expression of *Ceratophyllum demersum* phytochelatin synthase, CdPCS1, in *Escherichia coli* and *Arabidopsis* enhances heavy metal(loid)s accumulation. *Protoplasma* 250:1263–1272
- Siemianowski O, Barabasz A, Kendziorok M, Ruszczynska A, Bulska E, Williams LE, Antosiewicz DM (2014) AtHMA4 expression in tobacco reduces Cd accumulation due to the induction of the apoplastic barrier. *J Exp Bot* 65(4):1125–1139
- Singh NP, Santal AR (2015) Phytoremediation of heavy metals: the use of green approaches to clean the environment. In: Ansari AA, Gill SS, Gill R, Lanza GR, Newman L (eds) *Phytoremediation*. Springer, Cham, pp 115–129
- Song WY, Park J, Eisenach C, Maeshima M, Lee Y, Martinoia E (2014) ABC transporters and heavy metals. In: *Plant ABC transporters*. Springer International Publishing, New York, pp 1–17
- Taamalli M, Ghabriche R, Amari T, Mnasri M, Zolla L, Lutts S, Ghnaya T (2014) Comparative study of Cd tolerance and accumulation potential between *Cakilemaritima* L. (halophyte) and *Brassica juncea* L. *Ecol Eng* 71:623–627
- Takahashi R, Ishimaru Y, Shimo H, Bashir K, Senoura T, Sugimoto K et al (2014) From laboratory to field: *OsNRAMP5*-knockdown rice is a promising candidate for Cd phytoremediation in paddy fields. *PLoS One* 9(6):e98816
- Tiong JY, McDonald GK, Genc Y, Pedas P, Hayes JE, Toubia J, Huang CY (2014) HvZIP7 mediates zinc accumulation in barley (*Hordeum vulgare*) at moderately high zinc supply. *New Phytol* 201(1):131–143

- Tiong J, McDonald G, Genc Y, Shirley N, Langridge P, Huang CY (2015) Increased expression of six ZIP family genes by zinc (Zn) deficiency is associated with enhanced uptake and root-to-shoot translocation of Zn in barley (*Hordeum vulgare*). *New Phytol* 207(4):1097–1109
- Tripathi P, Singh PK, Mishra S, Gautam N, Dwivedi S, Chakrabarty D, Tripathi RD (2015) Recent advances in the expression and regulation of plant metallothioneins for metal homeostasis and tolerance. In: *Environmental waste management*. CRC Press, Boca Raton, pp 551–564
- Tsednee M, Yang SC, Lee DC, Yeh KC (2014) Root-secreted nicotianamine from *Arabidopsis halleri* facilitates zinc hypertolerance by regulating zinc bioavailability. *Plant Physiol* 166(2):839–852
- Vrbová M, Kotrba P, Horáček J, Smýkal P, Šváblová L, Vétrovcová M, Smýkalová I, Griga M (2015) Enhanced accumulation of cadmium in *Linum usitatissimum* L. plants due to overproduction of metallothionein a-domain as a fusion to b-glucuronidase protein. *Plant Cell Tissue Organ Cult* 112:321–330
- Wan X, Lei M, Chen T (2016) Cost–benefit calculation of phytoremediation technology for heavy-metal-contaminated soil. *Sci Total Environ* 563:796–802
- Wang QF, Zhao Y, Yi Q, Li KZ, Yu YX, Chen LM (2010) Overexpression of malate dehydrogenase in transgenic tobacco leaves: enhanced malate synthesis and augmented Al-resistance. *Acta Physiol Plant* 32(6):1209–1220
- Wang Q, Yi Q, Hu Q, Zhao Y, Nian H, Li K, Chen L (2012) Simultaneous overexpression of citrate synthase and phosphoenolpyruvate carboxylase in leaves augments citrate exclusion and Al resistance in transgenic tobacco. *Plant Mol Biol Rep* 30(4):992–1005
- Wang JW, Li Y, Zhang YX, Chai TY (2013) Molecular cloning and characterization of a *Brassica juncea* yellow stripe-like gene, BjYSL7, whose overexpression increases heavy metal tolerance of tobacco. *Plant Cell Rep* 32(5):651–662
- Wen XP, Ban Y, Inoue H, Matsuda N, Moriguchi T (2010) Spermidine levels are implicated in heavy metal tolerance in a spermidine synthase overexpressing transgenic European pear by exerting antioxidant activities. *Transgenic Res* 19(1):91–103
- Wen XP, Ban Y, Inoue H, Matsuda N, Kita M, Moriguchi T (2011) Antisense inhibition of a spermidine synthase gene highlights the role of polyamines for stress alleviation in pear shoots subjected to salinity and cadmium. *Environ Exp Bot* 72(2):157–166
- Williams LE, Pittman JK, Hall JL (2000) Emerging mechanisms for heavy metal transport in plants. *Biochim Biophys Acta Biomembr* 1465(1):104–126
- Wiszniewska A, Hanus-Fajerska E, Smoleń S, Muszyńska E (2015) In vitro selection for lead tolerance in shoot culture of *Daphne* species. *Acta Sci Pol Hort Cult* 14(1):129–142
- Xia Y, Qi Y, Yuan Y, Wang G, Cui J, Chen Y, Zhang H, Shen Z (2012) Overexpression of *Elsholtzia haichowensis* metallothionein 1 (EhMT1) in tobacco plants enhances copper tolerance and accumulation in root cytoplasm and decreases hydrogen peroxide production. *J Hazard Mater* 233–234:65–71
- Xue HB, Stumm W, Sigg L (1988) The binding of heavy metals to algal surfaces. *Wat Res* 22:917–926
- Yadav R, Arora P, Kumar S, Chaudhury A (2010) Prospectives for genetic engineering of population for enhanced phytoremediation abilities. *Ecotoxicology* 19(8):1574–1588
- Yang JL, Wang YC, Liu GF, Yang CP, Li CH (2011) Tamarixhispidametallothionein-like ThMT3, a reactive oxygen species scavenger, increases tolerance against Cd²⁺, Zn²⁺, Cu²⁺, and NaCl in transgenic yeast. *Mol Biol Rep* 38:1567–1574
- Yang JL, Chen Z, Wu S, Cui Y, Zhang L, Dong H, Li C (2015) Overexpression of the *Tamarix hispida* ThMT3 gene increases copper tolerance and adventitious root induction in *Salix matsudana* Koizd. *Plant Cell Tissue Organ Cult* 121(2):469–479
- Yazaki K, Yamanaka N, Masunto T, Konagai S, Kaneko S, Ueda K, Sato F (2006) Heterologous expression of a mammalian ABC transporter in plant and its application to phytoremediation. *Plant Mol Biol* 61:491–503
- Zdunić Z, Grljušić S, Ledenčan T, Duvnjak T, Šimić D (2014) Quantitative trait loci mapping of metal concentrations in leaves of the maize IBM population. *Hereditas* 151(2–3):55–60
- Zhang Y, Liu J (2011) Transgenic alfalfa plants co-expressing glutathione S-transferase (GST) and human CYP2E1 show enhanced resistance to mixed contaminants of heavy metals and organic pollutants. *J Hazard Mater* 189(1):357–362
- Zhang Y, Liu J, Zhou Y, Gong T, Wang J, Ge Y (2013) Enhanced phytoremediation of mixed heavy metal (mercury)–organic pollutants (trichloroethylene) with transgenic alfalfa co-expressing glutathione S-transferase and human P450 2E1. *J Hazard Mater* 260:1100–1107
- Zhang J, Zhang M, Tian S, Lu L, Shohag MJI et al (2014) Metallothionein 2 (SaMT2) from *Sedum alfredii* Hance confers increased Cd tolerance and accumulation in yeast and tobacco. *PLoS One* 9(7):e102750. <https://doi.org/10.1371/journal.pone.0102750>
- Zhao C, Xu J, Li Q, Li S, Wang P et al (2014) Cloning and characterization of a *Phragmites australis* phytochelatin synthase (PaPCS) and achieving Cd tolerance in tall fescue. *PLoS One* 9(8):e103771. <https://doi.org/10.1371/journal.pone.0103771>
- Zhou G, Pereira JF, Delhaize E, Zhou M, Magalhaes JV, Ryan PR (2014) Enhancing the aluminum tolerance of barley by expressing the citrate transporter genes SbMATE and FRD3. *J Exp Bot* 65(9):2381–2390