



## Review

## Phytoremediation: Environmentally sustainable way for reclamation of heavy metal polluted soils

Sana Ashraf<sup>a,\*</sup>, Qasim Ali<sup>a,b</sup>, Zahir Ahmad Zahir<sup>a</sup>, Sobia Ashraf<sup>a</sup>, Hafiz Naeem Asghar<sup>a</sup><sup>a</sup> Institute of Soil & Environmental Sciences, University of Agriculture Faisalabad, Pakistan<sup>b</sup> College of Agriculture, Bahauddin Zakariya University, Bahadur Sub-campus Layyah, Pakistan

## ARTICLE INFO

## Keywords:

Heavy metals  
Phytoremediation  
Phytoextraction  
Environmental sustainability  
Biological sulfur oxidation  
Acidified cow dung slurry

## ABSTRACT

Soil contamination with toxic metals is a widespread environmental issue resulting from global industrialization within the past few years. Therefore, decontamination of heavy metal contaminated soils is very important to reduce the associated risks and for maintenance of environmental health and ecological restoration. Conventional techniques for reclamation of such soils are expensive and environmental non-friendly. Phytoremediation is an emerging technology implementing green plants to clean up the environment from contaminants and has been considered as a cost-effective and non-invasive alternative to the conventional remediation approaches. There are different types of phytoremediation including, phytostabilization, phytostimulation, phytotransformation, phytofiltration and phytoextraction, the latter being most extensively acknowledged for remediation of soils contaminated with toxic heavy metals. Recent literature is gathered to critically review the sources, hazardous effects of toxic heavy metals and environmentally sustainable phytoremediation technique for heavy metal polluted soils to offer widespread applicability of this green technology. Different strategies to enhance the bioavailability of heavy metals in the soil are also discussed shortly. It can be concluded that phytoremediation of heavy metal contaminated soils is a reliable tool and necessary for making the land resource accessible for crop production.

## 1. Introduction

Heavy metal pollution of soil has become global environmental dilemma due to intensively increasing industrialization and agricultural activities (Chaoua et al., 2018; Woodford, 2019). Heavy metals are highly toxic because, unlike organic matter, they are not biodegradable but can only change their oxidation state and are highly persistent in nature with half-life more than 20 years (Jan et al., 2015; Ayangbenro, and Babalola, 2017; Hadia-e-Fatima and Ahmed, 2018). Fifty-three elements are documented as heavy metals and are considered as universal pollutants with densities greater than 5 g/cm<sup>3</sup> (Sarma, 2011; Saif and Khan, 2017; Prieto et al., 2018). Heavy metals can be grouped into essential and non-essential classes. Essential heavy metals include Co, Cr, Cu, Fe, Mn, Ni and Zn and are considered as essential micronutrients, but become poisonous when taken in excess quantities. Non-essential heavy metals include Pb, Cd and Hg and are highly toxic for living organisms (Monni et al., 2000; ul-Hassan et al., 2017; Sandeep et al., 2019).

Heavy metals are natural components of soil, but human activities have increased their concentration. Sources of heavy metals in soil

include excessive application of agrochemicals, sewage sludge, industrial wastewater, biosolids and manure (Bu-Olayan and Thomas, 2009; Alghobar and Suresha, 2017; Haroon et al., 2018; Woodford, 2019). Consequently, heavy metal accumulation in soil causes severe health problems for plants, animals and humans (Kleckerova and Docekalova, 2014; Srivastava et al., 2017). According to the United States Environmental Protection Agency (US EPA), soil heavy metal pollution has caused health issues for about 10 million humans all over the world (U.S. EPA, 2016). Hence, accumulation of heavy metals in crop plants through soil-root interface is a critical hazard (Maimon et al., 2009; Nazir et al., 2015; Sakizadeh and Ghorbani, 2017).

## 2. Sources of heavy metal pollution in soil

Anthropogenic and geological activities are two different sources through which heavy metals enter the soil (Keshavarzi et al., 2015; Liu et al., 2018). Industrial and domestic wastewater, smelting, mining, fuel manufacturing and agrochemicals are different anthropogenic input points of heavy metals in agricultural and non-agricultural soil (Wei and Yang, 2010; Marrugo-Negrete et al., 2017; Yan et al., 2018).

\* Corresponding author.

E-mail address: [sanaashrafenv@gmail.com](mailto:sanaashrafenv@gmail.com) (S. Ashraf).<https://doi.org/10.1016/j.ecoenv.2019.02.068>

Received 25 November 2018; Received in revised form 18 February 2019; Accepted 21 February 2019

Available online 14 March 2019

0147-6513/ © 2019 Elsevier Inc. All rights reserved.




Fig. 1. Sources of heavy metals in soil.

Heavy metals are released into atmosphere through discharge of gases and dust from transport and production of constructed material and energy. In atmosphere, heavy metals exist in the form of aerosols and then through precipitation become deposit to soil (Aksu, 2015; Naderizadeh et al., 2016; Suvarapu and Baek, 2017). Sources of heavy metals in soil are shown in Fig. 1. Excessive application of biosolids, fertilizers, pesticides and herbicides to crops causes buildup of these chemicals in soil and uptake by plant (Atafar et al., 2010; Bitew and Alemayehu, 2017). Application of different biosolids as valuable fertilizers such as composts, sewage sludge and livestock manure to agricultural land leads towards elevating the level of heavy metals in soil (Basta et al., 2005; Bitew and Alemayehu, 2017). Biosolids which are organic material also have different heavy metal contents like Pb, Cd, Hg and Cr. Heavy metals in biosolids applied to soil can also leach to groundwater (Brisolara and Qi, 2013). Concentration of some heavy metals in fertilizers and sewage sludge is shown in Table 1.

Micronutrient (Fe, Zn, Mn, Cu, Co and Ni) deficient soils are provided with these elements for healthy growth of plants and crops (Tripathi et al., 2015). Large amounts of fertilizers are being applied to agricultural soils to supply the crops with adequate quantity of nitrogen, phosphorus and potassium. The compounds which are used to provide these components also possess very minute quantities of toxic heavy metals. Phosphate fertilizers contain considerable amount of toxic heavy metals as compared to potash and nitrogen fertilizers. Due to excessive application of such fertilizers, concentration of heavy metals (Pb and Cd) is going to be increased (Boyd, 2010; Bitew and Alemayehu, 2017). Use of sewage water for irrigation also leads

towards contamination of soil and crops (Bridge, 2004; Chaoua et al., 2018; Haroon et al., 2018).

Pesticides which are being used in agriculture and horticulture sectors are composed of considerable amount of heavy metals such as copper sulfate and lead arsenate compounds are used in pesticides to control pests (McLaughlin et al., 2000; Missimer et al., 2018). Application of domestic and industrial wastewater in agriculture is a common way to solve irrigation problem in arable land. The level of heavy metals in sewage water is low but excessive and long-term irrigation of wastewater effluents has caused the buildup of heavy metals in soil where they fixed in different ways (Gupta et al., 2012; Chaoua et al., 2018). Extent of heavy metals in soil can range from traces to 100,000 mg/kg depending upon type and location of compound (Alloway, 2013; Liu et al., 2018). Heavy metals are main cause of unhealthy environmental condition because they can leach and are non-degradable (Martley et al., 2004; Alloway, 2013; Sidhu, 2016).

### 3. Environmentally toxic heavy metals and their classification

Heavy metals can be classified into essential and non-essential groups. Group of essential heavy metals comprises of Co, Cr, Cu, Fe, Mn, Ni and Zn and other includes Pb, Cd and Hg. Heavy metals can also be classified as extremely poisonous, moderately poisonous and relatively less poisonous according to their level of toxicity as shown in Table 2. Among non-essential heavy metals, Cd is more bioavailable and bioaccumulate due to its high solubility. Cd has no biological function and in compound form it is present as divalent ion (Volland et al., 2014; Oves et al., 2016). In heavy metal contaminated soil, Cd is mostly present with Zn that is an essential micronutrient. At high concentration Zn causes toxicity in plants, while Cd rarely causes phytotoxicity (Ryzhenko et al., 2017). Vetiver grass can be utilized for the reclamation of Cd contaminated soil because it can uptake and accumulate large concentration of Cd (Banerjee et al., 2016, 2019). In rhizosphere, microorganisms cause sequestration of Cd and thus influence the uptake of Cd by plants (Chen et al., 2019).

In roots of plants that are used for remediation of Cd contaminated soil, high concentration of glutathione and glutathione-S-transferase has found to be associated with detoxification processes (Gill et al., 2013; Wu et al., 2017). Among edible crops, highest level of Cd and Zn was found in maize and that of Pb was found in wheat. Use of Cd contaminated soil for wheat cultivation can cause food contamination because Cd uptake in wheat grains is high as compared to maize (Perrier et al., 2016). Cd has the potential to suppress the activities of antioxidative enzymes mainly glutathione reductase (Wu et al., 2017).

It is very difficult to remove Pb from contaminated soil. Soil's capacity to bind Pb improves with rising level of pH, cation exchange

Table 1

Concentration of heavy metals in fertilizer (Phosphate rock) and sewage sludge.

Source: (Edgell, 1988; Kongshaug et al., 1992; Nishimune, 1993; Wallace and Wallace, 1993; Benckiser and Simmarmata, 1994; Pantelica and Salagean, 1997; European Commission, 2001; Ogunleye et al., 2002; Conceicao and Bonotio, 2006; Jamali et al., 2007; Sabiha et al., 2009).

Country	Metal mg/kg dry material						
	As	Cd	Cr	Cu	Ni	Pb	Zn
<b>Phosphate rock</b>							
Brazil	–	4	70.5	–	116	44.5	299
Morocco	–	30	291	–	26	7	345
Nigeria	–	–	28	–	–	–	59
North Africa	–	60	105	–	33	6	420
Pakistan	–	7.5	17	–	28	89	67.2
Russia	–	0.1	23.3	–	2	3	19
USA	–	11	142	–	37	12	403
<b>Sewage Sludge</b>							
Germany	–	20–40	1000–1750	1000–1750	300–400	750–1200	2500–4000
Japan	75	85	3000	4300	420	840	7500
Pakistan	4.62	27.3	75.9	177.8	83.0	103.3	406.4
USA	–	10	900	800	100	900	2500

**Table 2**

Classification of heavy metals according to their level of toxicity.

Source: (GOST, 1983; Mukesh et al., 2008).

Heavy metal	Level of toxicity
Cadmium (Cd), Arsenic (As), Mercury (Hg), Selenium (Se), Lead (Pb), Zinc (Zn)	Extremely poisonous
Nickel (Ni), Cobalt (Co), Chromium (Cr), Copper (Cu), Molybdenum (Mo)	Moderately poisonous
Barium (Ba), Manganese (Mn), Strontium (Sr)	Relatively less poisonous

capacity, organic matter content and phosphate (Tang et al., 2015; Wijayawardena et al., 2015; Zeng et al., 2017). In soil, Pb becomes precipitated as Pb-phosphate and thus becomes unavailable to plants (Zeng et al., 2017). *Brassica juncea* can be utilized to extract Pb from contaminated soil because it transports Pb from root to shoot (Bouquet et al., 2017; Yahaghi et al., 2018). In fruit crops, Pb does not accumulate in their fruiting parts. High concentration of Pb has been found on the surface layer of leafy vegetables and root crops. Pb contaminated soil at a level more than 300 ppm affects the plant survival (Rosen, 2002; Chandrasekhar, and Ray, 2019).

Mercury (Hg) is the liquid metal and byproduct of ore processing. In environment, Hg exists in various forms. Mercuric and mercurous are oxidized forms, while organic and inorganic Hg become reduced to elemental Hg under reducing conditions that can be converted to alkylated form. Alkylated mercury is most hazardous, volatile and soluble in water (Smith et al., 1995; Sun et al., 2016; Beckers, and Rinklebe, 2017). Alkylated mercury is most toxic because due to its strong affinity for sulfur bounded compounds like enzymes and proteins.

Due to rising anthropogenic emissions, Zn has built up in the soil and water systems. Zn is phytotoxic and only some plants have ability to survive under Zn contaminated soil (Pence et al., 2000; Ryzhenko et al., 2017). Zn is a micronutrient and necessary for human health. Its deficiency can cause birth defects (Wang et al., 2015; Calderón Guzmán et al., 2019). Among heavy metals, cobalt (Co) is least toxic and forms different inorganic and organic compounds. In biological systems, Co works as a cofactor for different enzymes. However, at higher concentration it is hazardous for human health (Tchounwou et al., 2012; Leyssens et al., 2017). Nickel (Ni) is an essential element but at higher concentration it causes health effects. Nickel contaminated soil affects microbial growth and microorganisms also develop resistance against Ni (Wuana and Okieimen, 2011; Paulo et al., 2017; X. Xia et al., 2018).

Chromium (Cr) is an ore product of mining industry and in environment present as trivalent ( $\text{Cr}^{3+}$ ) and hexavalent ( $\text{Cr}^{6+}$ ) chromium (Smith et al., 1995; Mandal et al., 2017; Coetzee et al., 2018). Trivalent chromium is an essential trace element and cofactor for many enzymes (Davis and Vincent, 1997; Lewicki et al., 2014). Under aerobic conditions chromium exists as  $\text{Cr}^{6+}$  and under anaerobic conditions, soil organic matter converts  $\text{Cr}^{6+}$  into  $\text{Cr}^{3+}$ . In soils at low pH,  $\text{Cr}^{3+}$  is dominant that form complexes with  $\text{Cl}^-$ ,  $\text{CN}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{NH}_3$ ,  $\text{OH}^-$  (Chrostowski et al., 1991; Gautam et al., 2016).

Worldly, copper (Cu) is mostly used metal because it is an essential micronutrient for both plants and animals. In humans, Cu is important for hemoglobin production while in plants it is important for water regulation and seed production. Copper contaminated soil can cause direct and indirect threats on plants and humans, respectively (Bjuhr, 2007; Jaiswal et al., 2018). In soil, Cu forms complexes with organic matter. Thus, concentration of ionic Cu in soil solution is very low (Wuana and Okieimen, 2011; Lockwood et al., 2015). Copper is toxic due to its radical property that forms superoxide radicals which interact with thiol compounds in cell membrane (Nies, 2003).

Arsenic (As) is the most abundant metalloid in nature and found as arsenate ( $\text{As}^{5+}$ ) and arsenite ( $\text{As}^{3+}$ ) (Mukhopadhyay et al., 2002; Watanabe and Hirano, 2013). Arsenate is dominant under aerobic conditions while arsenite is dominant under anaerobic conditions. Arsenic forms methylated compounds that are volatile (Scragg, 2006; Schwartz et al., 2016). Arsenate behaves like phosphate and can enter

the microbial cells and prevent the phosphate dependent energy production processes and thus impede oxidative phosphorylation (Lloyd and Oremland, 2006; Urrialde et al., 2017).

#### 4. Heavy metal pollution and toxic effects on environment

Heavy metals are main source of environmental pollution because they can leach and are nondegradable (Martley et al., 2004; Brisolara and Qi, 2013; Sidhu, 2016; Chaoua et al., 2018; Woodford, 2019). Released heavy metals into soil have detrimental effects on living organisms through bioaccumulation and biomagnification processes (Szykowska et al., 2018; Mortensen et al., 2018). Most heavy metals are toxic for both micro and macro organisms by direct impacts on their physiological and biochemical pathways (Rai et al., 2016; Luo et al., 2019; Ghorri et al., 2019) and their concentration in soil ranges from 1 to 100,000 mg/kg (Blaylock and Huang, 2000; Herawati et al., 2000; Long et al., 2002; Alloway, 2013). The highest concentration of heavy metals occurs normally in the topsoil because organic horizons are characterized by the greatest ability to bind heavy metals (Acosta et al., 2015; Gu et al., 2016). Hence, high concentrations of Pb and Cd in upper layer of soil irrigated with contaminated water are taken up by plants (in rhizosphere, root exudates and microorganisms increase mobility of heavy metals) and thus negatively affect the absorption process of essential nutrients (Hédiji et al., 2015; Khan et al., 2016). Lead and cadmium can influence the absorption of  $\text{Cu}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Mn}^{2+}$  through competition for sites or processes shared by these cations (Xu and Shi, 2000). Hence, metal contaminated crops mainly vegetables are not safe for human edible purpose due to their network of roots in upper layer of soil (Smolen et al., 2010; Qureshi et al., 2016; Gupta et al., 2019). The maximum permissible limits of toxic heavy metals in irrigation water, soil and plants are presented in Table 3.

Heavy metal intoxicated plants in humans cause carcinogenic and chronic diseases. Human consumption of Pb, Cd and Zn causes respiratory diseases and heart, brain and kidney damage. Heavy metals cause toxicity in biological molecules in three different ways by producing reactive species, blocking functional groups and replacement of basic metal ions (Schutzendubel and Polle, 2002; Morcillo et al., 2016; Wijayawardena et al., 2016; Mandal, 2017). Heavy metals toxicity in plants leads towards stunted growth, low yield and aberrations in metabolic functions (Garbisu and Alkorta, 2001; Schwartz et al., 2003;

**Table 3**

Maximum permissible limits for toxic heavy metals concentration in irrigation water, soil and plants.

Source: (European Union, 2002; WHO/FAO, 2007).

Metal	Irrigation water ( $\mu\text{g}/\text{mL}$ )	Soil ( $\mu\text{g}/\text{g}$ )	Plant ( $\mu\text{g}/\text{g}$ )
Lead (Pb)	0.015	300	0.30
Cadmium (Cd)	0.01	3	0.2
Chromium (Cr)	0.10	150	5
Arsenic (As)	0.01	20	0.1
Nickel (Ni)	1.40	50	67
Mercury (Hg)	0.01	30	0.03
Copper (Cu)	0.20	140	40
Iron (Fe)	0.50	50,000	450
Zinc (Zn)	2.0	300	60
Manganese (Mn)	0.20	80	500

Asati et al., 2016; Kalaivanan and Ganeshamurthy, 2016). Increased production rate of reactive oxygen species (ROS) is the most obvious outcome of heavy metal toxicity in plants that causes oxidative stress leading towards membrane dismantling (Chen et al., 2012; Pandey and Dubey, 2019). Soil heavy metal contamination can also change native microbial community that ultimately destroys its biochemical properties (Kurek and Bollag, 2004; Paulo et al., 2017; X. Xia et al., 2018; B. Jiang et al., 2019; M. Jiang et al., 2019).

## 5. Remediation techniques for heavy metal-polluted soil

Decontamination of heavy metal contaminated soils is a main concern in environmental legislation. In-situ and ex-situ are the two approaches that are being practiced for decontamination of heavy metal affected soils. The ex-situ decontamination of polluted soils carried by conventional physico-chemical techniques is expensive. However, in situ remediation of heavy metal affected soils by plants is economical and environment friendly and depends upon the bioavailability of heavy metals (Van Gestel et al., 1992; Khalid et al., 2017; Emenike et al., 2018; Xia et al., 2019). Reclamation processes for heavy metal contaminated soils are classified into physical, chemical and biological processes (Zhou and Song, 2004; Khalid et al., 2017; Emenike et al., 2018). Different remediation techniques for heavy metal-polluted soil are shown in Fig. 2. Remediation mechanisms are based on two basic principles. First is the complete removal of contaminants and second is the transformation of contaminants into less harmful forms by using engineering technologies (Zhou and Song, 2004; Suthersan et al., 2016). Conventional physico-chemical soil remediation methods can be applied at massively heavy metal affected soils but only to small regions (Bio-Wise, 2003; Dada et al., 2015). Application of these methods is limited due to high energy requirement and adverse effects on soil structure and productivity (Leumann et al., 1995; Schnoor, 1997; Danh et al., 2009; Kuppusamy et al., 2017). Microorganisms can be applied for restoration of heavy metal affected soil (Garbisu and Alkorta, 1997; Ayangbenro, and Babalola, 2017). But use of plants for soil reclamation is more beneficial because they extract heavy metals from soil and restore the soil to healthy level (Chirakkara et al., 2016; Sarwar et al., 2017; Emenike et al., 2018; Lajayer et al., 2019).

### 5.1. Physico-chemical remediation techniques for heavy metal-contaminated soil

The physical methods of soil reclamation include soil replacement and thermal desorption processes. Soil replacement includes the use of contaminant-free soil to replace the polluted soil in order to dilute the

level of pollutants (Qian and Liu, 2000; Zhang et al., 2004; Fenyvesi et al., 2019). There are three types of soil replacement:

- i) Contaminated soil is removed and new soil is introduced
- ii) Contaminated soil is dugged up and pollutants are spread into deep layers
- iii) Imported new soil is added into contaminated soil (Sidhu, 2016; Khalid et al., 2017).

In thermal desorption contaminated soil is heated in order to volatile the pollutants. Contaminated soil is excavated and treated in thermal desorber. Contaminated vapors are captured by gas collection equipment. Thermal desorption is divided into low temperature and high temperature desorption (Aresta et al., 2008; Zhang et al., 2017). Chemical remediation technology of heavy metal contaminated soil includes immobilization techniques and soil washing. Immobilization techniques involve the application of organic and inorganic binding agents to contaminated soil for stabilization of contaminants by the natural processes of sorption, precipitation and complexation. In-situ immobilization processes are preferred due to low inputs of labor and energy (Hashimoto et al., 2009; Sidhu, 2016; Khalid et al., 2017).

#### 5.1.1. Solidification and stabilization

Solidification and stabilization are clean up technologies that avoid or slow the release of heavy metals or other contaminants. These methods prevent heavy metals from leaching by binding. Solidification involves the mixing of contaminated material with binding agent that causes the contaminated material to fix together (Evanko and Dzombak, 1997; Xia et al., 2017; W.Y. Xia et al., 2018). Inorganic binding agents include clay, zeolite, charcoal, Fe/Mn oxides, calcium carbonate, fly ash and cement (Fawzy, 2008; Gunatilake, 2015; Zarschler et al., 2016) and organic binding agents include manure, compost and bitumen (Farrell et al., 2010; Lim et al., 2016). Organic and inorganic binding agents can be used in a combination. The dominant mechanism of metal immobilization includes precipitation of hydroxides within solid medium. Solidification involves the mixing of contaminated material with binding agents but those binding agents also have chemical reactions with pollutants to prevent them from leaching into environment (Xia et al., 2017; W.Y. Xia et al., 2018).

#### 5.1.2. Vitrification

In vitrification, metal contaminated soil is heated at high temperature to reduce the mobility of heavy metals. During this process high temperature may cause volatilization of organic compounds and some metals such as Hg that should be collected and remaining are converted




Fig. 2. Different remediation techniques for heavy metal-polluted soil.

into inert glass like product (Asante-Duah, 1996; Kołacinski et al., 2017; RoyChowdhury et al., 2018). There are three major classes of vitrification process which include electrical, thermal and plasma methods (Acar and Alshawabkeh, 1993; Suthersan, 1997; Wait and Thomas, 2003). In-situ vitrification is preferred over ex-situ process and different energy sources can be applied for this technique (USEPA, 1992). In this process an electric current is passed through contaminated soil by using electrodes inserted into the soil. An array of four electrodes is called as a melt and a single melt can decontaminate up to 1000 t of polluted soil (Buelte and Thompson, 1992; Kołacinski et al., 2017).

### 5.1.3. Soil washing

Soil washing includes the implementation of solvents and mechanical methods to clean up heavy metal contaminated soil (Urum et al., 2003; Dermont et al., 2008; Song et al., 2017). In this technique, fine soil portion is separated from coarse soil that causes reduction in volume of contaminated soil (Riser-Roberts, 1998; Liao et al., 2016). The reduced contaminated soil volume can be treated by other clean up technologies while larger clean soil volume can be managed by utilizing as backfill (RAAG, 2000; Chu and Chan, 2003; Kumpiene et al., 2017). A range of contaminants in soil can be treated by soil washing including heavy metals (Park et al., 2002; Juhasz et al., 2003; Liao et al., 2016). Soil washing includes an in-situ flushing process with washing solvents followed by extraction of heavy metals by ex-situ process from soil slurry (Reed et al., 1995; Dermont et al., 2008). The effectiveness of this process depends upon the potential of washing solvents to dissolve the heavy metals present in the soil (Gombert, 1994; Dermont et al., 2008). Solvents are used according to the nature of contaminants present in the soil. A range of chemicals are applied as solvents for soil washing that includes organic acids, chelating compounds, surfactants and cosolvents (Gao et al., 2003; Maturi and Reddy, 2008; Zhang et al., 2009).

## 5.2. Biological remediation technique of heavy metal-contaminated soil: an environmentally sustainable way

Biological remediation approach that is an environmentally sustainable way to reclaim heavy metal contaminated soil includes bioremediation, phytoremediation and combination of both techniques.

### 5.2.1. Bioremediation

The principal objective of microbial reclamation of heavy metal polluted soil is to immobilize and reduce bioavailability of metals. Heavy metals cannot be degraded by microorganisms but can be converted to another form due to their altered physical and chemical properties. Efficiency of microbial remediation varies with the type of heavy metal and microorganism. Extracellular complexation, intracellular accumulation, oxidation-reduction reactions and precipitation are microbial remediation mechanisms. Microbial leaching is also a remediation tool for heavy metal contaminated soil (Q. Yang et al., 2016; Z. Yang et al., 2016; Yang et al., 2018). Biosorption is the most important mechanism in microbial remediation. Extracellular materials cause immobilization of heavy metals by binding with anionic functional groups of cell surface (Ayangbenro, and Babalola, 2017; Etesami, 2018; Gupta, and Diwan, 2017). Binding forces between metal ions and cell surfaces include covalent bonding, Vander Waals forces and electrostatic interactions (Blanco et al., 2000).

Siderophores that are chelating agents produced by plants, bacteria and fungi play a significant role in microbial remediation. Siderophores bind heavy metals and thus reduce their bioavailability (Schalk et al., 2011; Saha et al., 2016). Biosurfactants produced by microorganisms have the potential to complex heavy metals (Sarubbo et al., 2015; Ma et al., 2016). Complexation increases the dissolution of heavy metals that results in the reduction of metals and production of less soluble salts of metals. Microorganisms can remove heavy metals from environment through enzymatic or non-enzymatic processes (Nealson

et al., 1992; Sarubbo et al., 2015; Ma et al., 2016).

### 5.2.2. Phytoremediation

The objective of soil reclamation methods is to find a solution that is ecofriendly (Martin and Ruby, 2004). Phytoremediation is also called as agro-remediation, botano-remediation or green remediation. Phytoremediation is a green approach and sustainable way for soil reclamation as compared to conventional soil remediation techniques (Mahar et al., 2016). The concept of phytoremediation was first introduced in 1983 and still this technique is at testing stage. Phytoremediation is a best approach to deal with low to average metal contaminated soils and can be applied in combination with other traditional soil remediation approaches for efficient removal of contaminants (Cunningham and Ow, 1996; Helmisaari et al., 2007; Khalid et al., 2017).

Phytoremediation is an environment friendly mechanism in which fast growing plants are used to eliminate, hold or provide nontoxic contaminants in soil or water (Mahar et al., 2016). It is an aesthetical, economical and environment friendly way to detoxify contaminants (Zhang et al., 2010). Phytoremediation can be successful when used plant species can uptake and store high concentration of metal contaminants in their shoot parts. Phytoremediation technology is grouped into five sub classes including phytostabilization, phytostimulation, phytotransformation, phytofiltration and phytoextraction (Parmar, and Singh, 2015; Ramanjaneyulu et al., 2017).

**5.2.2.1. Phytostabilization.** Phytostabilization, phytoimmobilization or phytoreclamation is plant-based inactivation approach to deal with metal contaminated soil (Singh, 2012; Ramanjaneyulu et al., 2017). The objective of this technique is reduction in the mobility and bioavailability of heavy metals and consequently limits their leaching and entry into groundwater and food chain, respectively (Erakhrumen, 2007; Khalid et al., 2017). In phytostabilization plants play secondary role as compared to soil amendments. This technique involves physical and chemical immobilization of metal contaminants by their sorption onto roots and fixation with different soil amendments (Schnoor, 2000; Wuana and Okieimen, 2011). Organic matter, biosolids, clay minerals and phosphate fertilizers are most efficient soil amendments for immobilization of heavy metals (Flathman and Lanza, 1998). The main objectives of plants are to reduce water percolation, limit contact with contaminants, reduce soil erosion and decrease migration of contaminants (Raskin and Ensley, 2000; Akhtar et al., 2013). Phytostabilization is a management strategy and not a long-lasting way because at the end metal contaminants persist in the soil (Vangronsveld et al., 2009).

**5.2.2.2. Phytostimulation.** Phytostimulation also referred to as rhizodegradation is the disintegration of organic pollutants in rhizosphere with enhanced microbial activity (Mukhopadhyay and Maiti, 2010). Rhizosphere is soil volume about 1 mm nearby root and is influenced by root activity (Pilon-Smits, 2005). In rhizosphere, microbial activity is enhanced in different ways:

- i) Root exudates that contain amino acids and carbohydrates enrich indigenous microorganisms;
- ii) Roots ensure the supply of oxygen in rhizosphere for aerobic transformations;
- iii) Root biomass enhanced the availability of organic carbon;
- iv) Mycorrhizae fungi cause the degradation of compounds that cannot be breakdown by bacteria;
- v) Plants provide habitat for enhanced microbial population (Yadav et al., 2010).

**5.2.2.3. Phytotransformation.** Phytotransformation also termed as phytodegradation is the breakdown of organic compounds either by metabolic processes of plants or enzymes produced by plants and does not depend on microbial community (Vishnoi and Srivastava, 2008;

Kumar et al., 2018). Thus, plants can be considered as green liver of biosphere. Phytodegradation is only beneficial for breakdown of organic compounds. Release of volatile compounds to atmosphere through plant transpiration is termed as phytovolatilization. In this technique plants uptake contaminants from soil, transform them to volatile compounds and then discharge those volatile compounds into atmosphere. This technique is limited to organic compounds and Hg and Se heavy metals (Padmavathamma and Li, 2007; Kumar and Gunasundari, 2018).

**5.2.2.4. Phytofiltration.** Phytofiltration is the use of plant roots for reclamation of surface and groundwater and wastewater with low level of contaminants (Mukhopadhyay and Maiti, 2010). Firstly, plants are supplied with contaminated water to acclimate the plants and then these plants are transferred to contaminated site for remediation. When roots become saturated, they are harvested (Zhu et al., 1999). Phytofiltration can be rhizofiltration (using plant roots), blastofiltration (seedlings) or caulofiltration (plant shoots) (Mesjasz-Przybylowicz et al., 2004; Da Conceição Gomes et al., 2016). In this technique contaminants become absorbed, adsorbed or precipitated and their leaching to groundwater becomes reduced. Root exudates change rhizosphere pH and can cause precipitation of metals on plant roots (Flathman and Lanza, 1998; Javed et al., 2019). In rhizofiltration both terrestrial and aquatic fast-growing plants can be implemented for the extraction of Cd, Cr, Cu, Ni, Pb and Zn. Mostly terrestrial plants are used due to their fibrous and longer root system (Dhanwal et al., 2017).

**5.2.2.5. Phytoextraction.** In phytoextraction, fast growing plants are implemented to remove heavy metals from the environment (soil and water) (Yanai et al., 2006; Van Nevel et al., 2007; Pajevic et al., 2016). Phytoextraction involves two approaches that include continuous or natural and chemically induced phytoextraction (Ghosh and Singh, 2005). Continuous phytoextraction is an approach which includes the removal of heavy metals by network of roots and then directed to upper plant tissues above the ground (Jadia and Fulekar, 2008) as shown in Fig. 3. Harvested plant biomass can be used for biogas production and can also be combusted. Combusted plant biomass can be used for metal recovery, fixed in bricks or disposed in deserted lands. This technique is also known as phytomining or biomining (Singh and Bhargava, 2017a, 2017b). This approach is best to reduce concentration of metal contaminants from soil by plant roots and shoots without affecting soil properties. Metals can be restored from harvestable plant parts (Garbisu and Alkorta, 2001; Singh and Bhargava, 2017a, 2017b). Continuous cropping and harvesting system can cut down level of contaminants in soil (Vandenhove et al., 2001). In contrast to prevailing remediation techniques, this emerging green technology would be ten

times more economical (Wan et al., 2016).

Plants use for phytoextraction should have rapid growth, high biomass and extended roots network and should bear and store high levels of heavy metals (Romkens et al., 2002; Sytar et al., 2016). In continuous phytoextraction natural hyperaccumulators are used. Hyperaccumulator plants are used at metalliferous sites (Lasat, 2002; Ghosh and Singh, 2005; Usman et al., 2018). Phytoextraction involves different processes such as

- (i) Some metal fraction becomes sorbed at root surface
- (ii) Bioavailable metals enter the roots through cellular membrane
- (iii) Small fraction of metals taken up by roots becomes immobilized in vacuole
- (iv) Mobile metals in roots enter the xylem
- (v) Metals are translocated from roots to tissues of stems and leaves (Lasat, 2000; Adrees et al., 2015).

## 6. Use of metallophytes for phytoextraction

Plants that can survive in heavy metal contaminated soil are called as metallophytes (Sheoran et al., 2011). Metallophytes are mostly belonged to Brassicaceae plant family and are categorized into three classes; excluders, indicators and hyperaccumulators (McGrath et al., 2002; Bothe, 2011). Metal excluders uptake heavy metals from environment and accumulate into roots but limit their transport towards above ground plant tissues (Malik and Biswas, 2012). Metal indicators uptake contaminants from contaminated soil and accumulate into their upper parts (Sheoran et al., 2011). Metal hyper accumulator should accumulate at the minimum 100 mg/kg As and Cd, 1000 mg/kg Co, Cu, Cr, Ni and Pb and 1000 mg/kg Mn and Ni (Watanebe, 1997; Reeves and Baker, 2000). Heavy metals hyperaccumulation behaves as a defense action in response to plant infective agents (Boyd et al., 1994). Above 400 plant varieties have been recognized as hyperaccumulators with sluggish growth and less biomass production (McGrath and Zhao, 2003; Kramer, 2018). Field implementation of natural hyperaccumulators can be carried out for both phytoremediation and phytomining of heavy metals and precious metals, respectively (Brooks et al., 1998). Some important hyperaccumulators for heavy metal remediation are presented Table 4.

## 7. Impediments in the success of phytoextraction

The success of phytoextraction depends upon the accessibility of heavy metals to selected plant roots that is directly related to solubility of metal contaminants in soil solution (Felix, 1997; Antoniadis et al., 2017). Hence, bioavailability is the limiting factor for phytoextraction. The dissolution of heavy metals in soil solution is primarily regulated




Fig. 3. Phytoextraction process including different management options of end product.

**Table 4**  
Important hyperaccumulators for heavy metals remediation.

Heavy metal	Plant species	Maximum reported concentration (mg/kg)	Reference
Cd	<i>Azollapinnata</i> , <i>Arabispinniculata</i>	740, 1127	(Rai, 2008; Zeng et al., 2009)
Cu	<i>Ipmoea alpine</i> , <i>Eleocharis acicularis</i>	12,300, 20,200	(Baker and Walker, 1989; Sakakibara et al., 2011)
Pb	<i>Euphorbia cheiraadenia</i>	1138	(Chehregani and Malayeri, 2007)
Ni	<i>Alyssum corsicum</i> , <i>Alyssum markgrafii</i>	18,100, 19,100	(Li et al., 2003; Bani et al., 2010)
Zn	<i>Thlapsicaeruleaescens</i> , <i>Potentilla griffithii</i>	19,410, 19,600	(Banasova and Horak, 2008; Hu et al., 2009)
As	<i>Pteris vittata</i> , <i>Pteris ryukyuensis</i>	23,000, 3647	(Dong, 2005; Srivastava et al., 2006)
Cr	<i>Phragmites australis</i> , <i>Pteris vittata</i>	4825, 20,675	(Calheiros et al., 2008; Kalve et al., 2011)
Mn	<i>Schimasuperba</i>	62,412.3	(Yang et al., 2008)

by pH (Baker and Walker, 1989; McNeil and Waring, 1992; Henry, 2000; Javed et al., 2019). High soil pH leads towards greater retention and low dissolution of metals in soil solution, cation exchange capacity and organic matter content (Bliefert, 1994; Li and Shuman, 1996; Tang et al., 2015; Wijayawardena et al., 2015; Zeng et al., 2017). Heavy metals in soil can be categorized into available, exchangeable and unavailable fractions depending upon their uptake by plants (Zhou and Song, 2004; Q. Wei et al., 2008). The successful phytoextraction requires the application of plant species having excessive biomass production, higher growth rate and intense potential to accumulate poisonous heavy metals in their above ground parts (Clemens et al., 2002; Odoemelam and Ukpe, 2008; Kramer, 2018).

## 8. Enhancement of phytoextraction efficiency

Phytoextraction efficiency can be enhanced by using the following approaches:

- Increasing heavy metal bioavailability in soil
- Increasing plant biomass and decreasing phytoextraction cycle

### 8.1. Common approaches to increase heavy metal bioavailability

Two common practices which are being implemented to enhance the phytoavailability of heavy metals include the use of synthetic chelates (Tahmasbian and Sinegani, 2016; Chen et al., 2017; Chhajro et al., 2018) and reduction in soil pH (Zhu et al., 2016a, 2016b; Beiyuan et al., 2017; Agrawal and Singh, 2018). The reduction in soil pH can be accomplished by utilization of acids or acid-producing fertilizers (Murtaza et al., 2015; Zhu et al., 2016a, 2016b; Zhao et al., 2016; Agrawal and Singh, 2018). Synthetic chelates such as EDTA enters the plant roots and form soluble complexes with metals which leads towards enhancing the accessibility of heavy metals (Wenzel et al., 2003; B. Jiang et al., 2019; M. Jiang et al., 2019). But these methods have hazardous effects on physical and bio-chemical features of soil (Singh Brar et al., 2015) and can also cause groundwater pollution (Aziz et al., 2015; Q. Yang et al., 2016; Z. Yang et al., 2016; Postigo et al., 2017). A sustainable way to enhance heavy metal bioavailability by lowering soil pH is the application of acidified manure (Ashraf, 2017).

#### 8.1.1. Chelate-assisted or induced phytoextraction

Due to slow growth rate and low biomass of plants, chemically induced phytoextraction has been proposed. In this approach high biomass and fast-growing crops are used to extract huge concentration of heavy metals whose mobility in soil is raised by chelating agents (Smolinska, 2015; Mani et al., 2015; Patra et al., 2018). Efficiency of phytoextraction is restricted by phytoavailability of heavy metals (Felix, 1997). Accumulation of heavy metals can be enhanced in huge biomass production plants for the development of phytoremediation. Heavy metal accumulation potential of non-hyperaccumulating plants can be enhanced by using chelating substances. Chelating agents have been isolated from plants that are used for extraction and detoxification of heavy metals (Saifullah et al., 2009; Wiszniewska et al., 2016).

Enhanced availability and extraction of metal contaminants from soil by plants treated with chelating agents can be linked to different aspects such as (i) an crease in concentration of heavy metals in soil solution (ii) enhanced movement of metal-EDTA complexes towards roots (iii) less binding of formed complexes with negatively charged cell components of plant cell wall (iv) destruction of physiological barriers in roots due to high level of resultant complexes (v) increased mobility of complexes as compared to free ions results in more translocation of metals from roots to shoots (Evangelou et al., 2007). To enhance the potential of phytoextraction process, EDTA (ethylene diamine tetra acetic acid) was used as a chelating compound in early 1990s. EDTA can enhance the accumulation of heavy metals up to 100-fold (Grcman et al., 2001; Ali and Chaudhury, 2016). Both EDTA and EDTA-heavy metal complexes are highly hazardous for plants and soil microbial community (Chen and Cutright, 2001; Vassilev et al., 2004). Biodegradability of EDTA is very low and it also increases the leaching of heavy metals and thus cause ground water pollution (Lu et al., 2017; Qiao et al., 2017).


Ethylene diamine disuccinate (EDDS) is naturally produced by microorganisms and is effective for enhancing uptake of heavy metals (Sidhu et al., 2018). Bioavailability of Cu, Ni and Zn is enhanced more by the application of EDDS as compared to EDTA while that of Cd and Pb is enhanced more by EDTA than EDDS (Meers et al., 2005; Song et al., 2016). Heavy metal-EDDS complexes enter the roots and then transported to shoots (Luo et al., 2005). EDDS is toxic to some plants but not to soil microorganisms. Nitrilotriacetic acid (NTA) is a biodegradable chelating substance and has not phytotoxic effects (Wenger et al., 2003; De Souza Freitas and Do Nascimento, 2009). NTA has been used to enhance the proficiency of phytoextraction by enhancing the bioavailability of heavy metals. After phytoextraction, harvested biomass can be managed by using for phytomining to produce bio-ore (McGrath and Zhao, 2003; Singh and Bhargava, 2017a, 2017b).

#### 8.1.2. Biological sulfur oxidation to reduce pH and enhance metal bioavailability in soil

Positive effect of elemental sulphur on soil metal solubility because of soil pH reduction has been well documented (Ye et al., 2010), as it possesses slow release acidifying characteristic and is readily available (Chien et al., 2011). Application of elemental sulfur in soil has been recommended to reduce pH and enhance dissolution and bioavailability of heavy metals in soil (Tichy et al., 1997; Seidel et al., 1998; Kayser et al., 2000; Li et al., 2017; Sun et al., 2018). *Thiobacillus* bacteria are the most active sulfur oxidizing bacteria in soil (Besharati, 2003; Zhao et al., 2017). Sulfur is the most common and cost effective natural acidifying element. In soil, *Thiobacillus* bacteria oxidize one mole of elemental sulfur and produce two moles of hydrogen ions in soil and thus cause acidification of soil (Kaplan and Orman, 1998). Karimizarchi et al (2016) incubated the soil for 40 days with elemental sulphur at the rates of 0.5, 1 and 2 g kg<sup>-1</sup> soil and observed that pH decreased from 7.51 to 6.66, 5.45 and 4.8, respectively. Soil temperature and moisture level are two important factors that influence the rate of microbial sulfur oxidation in soil. Production of acid in soil by *Thiobacillus* bacteria is also dependent on sulfur to soil total solids ratio (Tsai et al., 2003).


### 8.1.3. Novel approach to enhance metal bioavailability by reducing soil pH through the application of microbially augmented acidified cow dung

A novel approach to enhance the bioavailability of the heavy metals has been recently reported by Ashraf (2017) and Ashraf et al. (2018). Authors have reported an increase of 114% and 126% in Pb and Cd concentration, respectively, in the shoots of rye grass that has been achieved by application of the acidified product. The bioaugmented acidified cow dung has been applied to soil to enhance the bioavailability of Pb and Cd to grasses used for phytoextraction in pot trials. Cow dung was acidified by adding elemental sulfur (S<sup>0</sup>) and molasses solution and also bioaugmented with isolated heavy metal resistant sulfur oxidizing bacteria (SOB). Mechanism of cow dung acidification is shown by equation below (Ashraf, 2017; Ashraf et al., 2018).



Minimum inhibitory concentration (MIC) of added SOB for Pb and Cd was 1000 mg/L and 180 mg/L, respectively (Ashraf et al., 2018). SOB oxidized (S<sup>0</sup>) and produced sulfuric acid (H<sub>2</sub>SO<sub>4</sub>). Acidified cow dung was applied in slurry form at field capacity level for multiple times to grasses during phytoextraction period. Acidified cow dung enhanced the bioavailability of Pb and Cd by reducing the pH of soil. Diluted acidified cow dung reduced the soil pH by 0.92 point and enhanced the concentration of Pb and Cd in shoot of rye grass from 44 mg/kg to 94.32 mg/kg and 34 mg/kg to 77 mg/kg, respectively (Ashraf, 2017). So, (S<sup>0</sup>), molasses and SOB can be applied to cow dung to acidify the manure that can be used for phytoextraction of heavy metals. Further, composts prepared with a wide range of organic wastes have many useful applications such as soil restoration (Ashraf, 2017).

**8.1.3.1. Mechanism how acidified cow dung enhance the bioavailability of heavy metals (Pb and Cd).** Acidified cow dung is enriched with nutrients and microorganisms and also of low pH. Both nutrients and microorganisms enhanced plant growth and promoted phytoextraction because for phytoextraction, plant biomass should be high. Acidified cow dung reduced the soil pH for short time period and enhanced the bioavailability of Pb and Cd. Most of the Pakistani soil is calcareous in nature having pH 8 and above. At this pH metals become immobilized due to their absorption on CaCO<sub>3</sub> particles. However, metals become available at pH 7. Sulfate ions present in the acidified cow dung react with water and form sulfuric acid (H<sub>2</sub>SO<sub>4</sub>). When acidulated cow dung is applied, H<sub>2</sub>SO<sub>4</sub> reacted with CaCO<sub>3</sub> and dissolve it. As a result, CaSO<sub>4</sub> was formed that is a fertilizer as shown in equation below. Heavy metal cations released into soil solution taken up by plants. Further, acidified cow dung is a nutrient rich material. So that, cations present in acidified cow dung exchanged the metal cations present on soil colloidal particles such as clay minerals (Ashraf, 2017).



### 8.2. Increasing plant biomass and decreasing phytoextraction cycle

Plant biomass plays very significant function in the removal of heavy metals through phytoextraction. So, the potential of phytoremediation can be improved by the application of fertilizers and proper irrigation system (Jankong et al., 2007; S.H. Wei et al., 2008; Nie et al., 2010). Efficiency of phytoremediation can also be improved by decreasing phytoremediation cycle. Phytoremediation cycle can be decreased by providing specific requirements of used plant species. This objective can also be obtained by shifting the seedlings of specific plant species to the field in order to limit the time period of phytoremediation. This approach can be beneficial because accumulation of heavy metals in shoots of plants is maximum at flowering stage (Wu et al., 2009).

## 9. Quantification of phytoextraction efficiency

Potential of phytoextraction can be quantitatively assessed by using bioconcentration factor, translocation factor and time of phytoremediation (Zhuang et al., 2005). Bioconcentration factor is a measure of plant's efficiency for concentrating metals into its tissues from the surrounding environment (Ladislas et al., 2012). Translocation factor is a measure of plants's efficiency for translocating the concentrated metals from its roots to shoots (Padmavathamma and Li, 2007).

Thus, translocation factor is the ratio of concentration of heavy metals in above ground plant tissues to that in its roots (Mattina et al., 2003; Liu et al., 2010). For the selection of natural hyperaccumulators for phytoextraction of metals, both bioconcentration and translocation factors are important (Wu et al., 2011). Translocation factor for natural hyperaccumulators should be > 1 to show that the level of metals is greater in above ground tissues than below ground tissues. Thus, translocation factor is most important for phytoextraction which involves the harvesting of aerial parts of plants (Wei and Zhou, 2004; Karami and Shamsuddin, 2010).

## 10. Advantages, disadvantages and future trends in phytoremediation

Phytoremediation is the cheapest technology to treat heavy metal contaminated soil (Rakhshaei et al., 2009). This emerging technology is about 60–80% inexpensive as compared to conventional physico-chemical techniques (Mwegoha, 2008). The harvested plant biomass after phytoextraction can be used for bioenergy production (Van Nevel et al., 2007; Tian and Zhang, 2016; Tripathi et al., 2016; Dhiman et al., 2016). Phytoremediation is a promising clean up technology for heavy metal polluted soil. But, this technique also has some disadvantages like slow growth rate of metal accumulating plant species, low bioavailability of heavy metals and long time period of phytoremediation (Nees et al., 2011; Ramamurthy and Memarian, 2012).

Phytoremediation is an emerging field of research and is limited to pot level with few field studies. Results can be different under greenhouse and field conditions (Ji et al., 2011). Phytoremediation process at field level is affected by different factors such as uneven distribution of pollutants, soil pH, pathogens, nutrients, moisture and temperature. For commercialization this technology must be tested at field level (Vangronsveld et al., 2009). For the success of this technology, identification of plants species having high biomass production and heavy metals accumulation capacity is required (Rodriguez et al., 2005).

## 11. Conclusion

This review revealed that heavy metals are amongst the most critical threats to the soil and human health. These metals are released into the environment through different sources. Conventional remediation techniques are costly and environmentally devastating. Therefore, it is unavoidable to implement low cost and eco-friendly technologies to remediate heavy metal polluted soils. Phytoremediation of metals is the most effective plant-based approach to remove pollutants from contaminated areas without posing any destructive effects to soil structure. Some plants have potential to absorb toxic metals up to several percent of their dried shoot biomass, these plants are known as hyperaccumulators. New plant species suitable for removing heavy metals from contaminated soils should be found. Metal bioavailability enhancing strategy such as use of bioaugmented acidified manure can be applied to polluted soil to decrease the soil pH and thus to enhance the availability of heavy metals to plants.

## Conflicts of interest

The authors declare no conflict of interest.

## References

- Acar, Y.B., Alshawabkeh, A., 1993. Principle of electrokinetic remediation. *J. Environ. Sci. Technol.* 27 (13), 2638–2647.
- Acosta, J.A., Gabarron, M., Faz, A., Martínez-Martínez, S., Zornoza, R., Arocena, J.M., 2015. Influence of population density on the concentration and speciation of metals in the soil and street dust from urban areas. *Chemosphere* 134, 328–337.
- Adrees, M., Ali, S., Rizwan, M., Zia-ur-Rehman, M., Ibrahim, M., Abbas, F., Farid, M., Qayyum, M.F., Irshad, M.K., 2015. Mechanisms of silicon-mediated alleviation of heavy metal toxicity in plants: a review. *Ecotoxicol. Environ. Saf.* 119, 186–197.
- Agrawal, M., Singh, A., 2018. Reduction in metal toxicity by applying different soil amendments in agricultural field and its consequent effects on characteristics of radish plants (*Raphanus sativus* L.). *J. Agr. Sci. Technol.* 15, 1553–1564.
- Akhtar, M.S., Chali, B., Azam, T., 2013. Bioremediation of arsenic and lead by plants and microbes from contaminated soil. *Res. Plant Sci.* 1 (3), 68–73.
- Aksu, A., 2015. Sources of metal pollution in the urban atmosphere (A case study: tuzla, Istanbul). *J. Environ. Health Sci. Eng.* 13 (1), 1–10.
- Alghobhar, M.A., Suresha, S., 2017. Evaluation of metal accumulation in soil and tomatoes irrigated with sewage water from Mysore city, Karnataka, India. *J. Saudi Soc. Agric. Sci.* 16 (1), 49–59.
- Ali, S.Y., Chaudhury, S., 2016. EDTA-enhanced phytoextraction by *Tagetes* sp. and effect on bioconcentration and translocation of heavy metals. *Environ. Process.* 3 (4), 735–746.
- Alloway, B.J., 2013. Sources of heavy metals and metalloids in soils. In: Alloway, B.J. (Ed.), *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability*. Springer Science & Business Media, Dordrecht, pp. 11–50.
- Antoniadis, V., Levizou, E., Shaheen, S.M., Ok, Y.S., Sebastian, A., Baum, C., Prasad, M.N., Wenzel, W.W., Rinklebe, J., 2017. Trace elements in the soil-plant interface: phytoavailability, translocation, and phytoremediation-a review. *Earth Sci. Rev.* 171, 621–645.
- Aresta, M., Dibenedetto, A., Fragale, C., 2008. Thermal desorption of polychlorobiphenyls from contaminated soils and their hydrodechlorination using Pd- and Rh-supported catalysts. *Chemosphere* 70 (6), 1052–1058.
- Asante-Duah, D.K., 1996. *Managing Contaminated Sites: Problem Diagnosis and Development of Site Restoration*. Wiley, New York, NY.
- Asati, A., Pichhode, M., Nikhil, K., 2016. Effect of heavy metals on plants: an overview. *Int. J. Appl. Innov. Eng. Manag.* 5, 2319–4847.
- Ashraf, S., 2017. *Phytoextraction of Lead and Cadmium by Grasses from Contaminated Soil Amended with Acidulated Cow Dung Slurry/extract and Bioaugmented with Sulfur Oxidizing Bacteria*. (Ph.D. diss). Inst. Soil Env. Sci., Univ. Agric., Faisalabad, Pakistan.
- Ashraf, S., Zahir, Z.A., Asghar, H.N., Asghar, M., 2018. Isolation, screening and identification of lead and cadmium resistant sulfur oxidizing bacteria. *Pak. J. Agric. Sci.* 55 (2), 349–359.
- Atafar, Z., Mesdaghinia, A., Nouri, J., Homae, M., Yunesian, M., Ahmadi Moghaddam, M., Mahvi, A.H., 2010. Effect of fertilizer application on soil heavy metal concentration. *Environ. Monit. Assess.* 160 (1–4), 83–89.
- Ayangbenro, A.S., Babalola, O.O., 2017. A new strategy for heavy metal polluted environments: a review of microbial biosorbents. *Int. J. Environ. Res. Public Health* 14 (1), 94. <https://doi.org/10.3390/ijerph14010094>.
- Aziz, T., Maqsood, M.A., Kanwal, S., Hussain, S., Ahmad, H.R., Sabir, M., 2015. Fertilizers and environment: issues and challenges. In: Rehman, H.K. (Ed.), *Crop Production and Global Environmental Issues*. Springer, Cham, pp. 575–598.
- Baker, A.J.M., Walker, P.L., 1989. Ecophysiology of metal uptake by tolerant plants. In: Shaw, A.J. (Ed.), *Heavy Metal Tolerance in Plants: Evolutionary Aspects*. CRC, Boca Raton, FL, pp. 155–177.
- Banasova, V., Horak, O., 2008. Heavy metal content in *Thlaspi caerulescens*, J. et C. Presl growing on metalliferous and non-metalliferous soils in Central Slovakia. *Int. J. Environ. Pollut.* 33, 133–145.
- Banerjee, R., Goswami, P., Lavana, S., Mukherjee, A., Lavana, U.C., 2019. Vetiver grass is a potential candidate for phytoremediation of iron ore mine spoil dumps. *Ecol. Eng.* <https://doi.org/10.1016/j.ecoeng.2018.10.012>.
- Banerjee, R., Goswami, P., Pathak, K., Mukherjee, A., 2016. Vetiver grass: an environment clean-up tool for heavy metal contaminated iron ore mine-soil. *Ecol. Eng.* 90, 25–34.
- Bani, A., Pavlova, D., Echevarria, G., Mullaj, A., Reeves, R.D., Morel, J.L., Sulce, S., 2010. Nickel hyperaccumulation by the species of *Alyssum* and *Thlaspi* (Brassicaceae) from the ultramafic soils of the Balkans. *Bot. Serb.* 34, 3–14.
- Basta, N.T., Ryan, J.A., Chaney, R.L., 2005. Trace element chemistry in residual-treated soil: key concepts and metal bioavailability. *J. Environ. Qual.* 34 (1), 49–63.
- Beckers, F., Rinklebe, J., 2017. Cycling of mercury in the environment: sources, fate, and human health implications: a review. *Crit. Rev. Environ. Sci. Technol.* 47 (9), 693–794.
- Beiyuan, J., Awad, Y.M., Beckers, F., Tsang, D.C., Ok, Y.S., Rinklebe, J., 2017. Mobility and phytoavailability of As and Pb in a contaminated soil using pine sawdust biochar under systematic change of redox conditions. *Chemosphere* 178, 110–118.
- Benckiser, T., Simmarmata, T., 1994. Environmental impact of fertilizing soils by using sewage sludge and animal waste. *Fert. Res.* 37, 1–22.
- Besharati, H., 2003. Preparation of suitable preservative for bacteria genera *Thiobacillus* and study its interactions with mycorrhiza fungi. *Environ. Microbiol.* 3 (2), 31–39.
- Bio-Wise, 2003. *Contaminated Land Remediation: A Review of Biological Technology*. DTI, London.
- Bitew, Y., Alemayehu, M., 2017. Impact of crop production inputs on soil health: a review. *Asian J. Plant. Sci.* 16, 109–131.
- Bjuhr, J., 2007. *Trace Metals in Soils Irrigated with Waste Water in a Periurban Area downstream Hanoi City, Vietnam*. Ph.D. diss. Swedish University of Agricultural Sciences.
- Blanco, A., Sanpedro, M.A., Sanz, B., Llama, M.J., Serra, J.L., 2000. Immobilization of nonviable cyanobacteria and their use for heavy metal adsorption from water. In: Oluquin, E.J., Sanehez, G., Hernandez, E. (Eds.), *Environmental Biotechnology and Cleaner Bioprocesses*. Philadelphia Taylor and Francis, pp. 135–151.
- Blaylock, M.J., Huang, L.W., 2000. Phytoextraction of metals. In: Raskin, I., Ensley, B.D. (Eds.), *Phytoremediation of Toxic Metals: Using Plants to Clean-up the Environment*. John Wiley and Sons, New York, pp. 53–70.
- Briefert, C., 1994. *Umweltchemie*. Wiley Europe-VCH, Weinheim, New York.
- Bothe, H., 2011. *Plants in heavy metal soils*. In: Sherameti, I., Varma, A. (Eds.), *Detoxification of Heavy Metals*. Springer, Berlin, Heidelberg, pp. 35–57.
- Bouquet, D., Braud, A., Lebeau, T., 2017. *Brassica juncea* tested on urban soils moderately contaminated by lead: origin of contamination and effect of chelates. *Int. J. Phytoremediat.* 19 (5), 425–430.
- Boyd, R.S., 2010. Heavy metal pollutants and chemical ecology: exploring new frontiers. *J. Chem. Ecol.* 36, 46–58.
- Boyd, R.S., Shaw, J.J., Martens, S.N., 1994. Nickel hyperaccumulation defends *Streptanthus polygaloides* (Brassicaceae) against pathogens. *Am. J. Bot.* 81, 294–300.
- Bridge, G., 2004. Contested terrain: mining and the environment. *Annu. Rev. Environ. Resour.* 29, 205–259.
- Brisolara, K.F., Qi, Y., 2013. Biosolids and sludge management. *Water Environ. Res.* 85 (10), 1283–1297.
- Brooks, R.R., Chambers, M.F., Nicks, L.J., Robinson, B.H., 1998. Phytomining. *Trends Plant Sci.* 3, 359–362.
- Buelt, J.L., Thompson, L.E., 1992. *The in Situ Vitrification Integrated Program: Focusing on an Innovative Solution on Environmental Restoration Needs*. Battelle Pacific Northwest Laboratory, Richland, Wash, USA.
- Bu-Olayan, A.H., Thomas, B.V., 2009. Translocation and bioaccumulation of trace metals in desert plants of Kuwait Governorates. *Res. J. Environ. Sci.* 3 (5) (581–58).
- Calderón Guzmán, D., Juárez Olguín, H., Osnaya Brizuela, N., Hernández García, E., Lindoro Silva, M., 2019. The use of trace and essential elements in common clinical disorders: roles in assessment of health and oxidative stress status. *Nutr. Cancer* 19, 1–8.
- Calheiros, C.S.C., Rangel, A.O.S.S., Castro, P.M.L., 2008. The effects of tannery wastewater on the development of different plant species and chromium accumulation in *Phragmites australis*. *Arch. Environ. Contam. Toxicol.* 55, 404–414.
- Chandrasekhar, C., Ray, J.G., 2019. Lead accumulation, growth responses and biochemical changes of three plant species exposed to soil amended with different concentrations of lead nitrate. *Ecotoxicol. Environ. Saf.* 171, 26–36.
- Chaoua, S., Boussaa, S., El Gharmali, A., Boumezzough, A., 2018. Impact of irrigation with wastewater on accumulation of heavy metals in soil and crops in the region of Marrakech in Morocco. *J. Saudi Soc. Agric. Sci.* <https://doi.org/10.1016/j.jssas.2018.02.003>. <https://www.sciencedirect.com/science/article/pii/S1658077X17303521>.
- Chehregani, A., Malayeri, B.E., 2007. Removal of heavy metals by native accumulator plants. *Int. J. Agri. Biol.* 9, 462–465.
- Chen, H., Cutright, T., 2001. EDTA and HEDTA effects on Cd, Cr, and Ni uptake by *Helianthus annuus*. *Chemosphere* 45, 21–28.
- Chen, F., Gao, J., Zhou, Q., 2012. Toxicity assessment of simulated urban runoff containing polycyclic musks and cadmium in *Carassius auratus* using oxidative stress biomarkers. *Environ. Pollut.* 162, 91–97.
- Chen, Z., Tang, Y.T., Zhou, C., Xie, S.T., Xiao, S., Baker, A.J., Qiu, R.L., 2017. Mechanisms of Fe biofortification and mitigation of Cd accumulation in rice (*Oryza sativa* L.) grown hydroponically with Fe chelate fertilization. *Chemosphere* 175, 275–285.
- Chen, X.W., Wu, L., Luo, N., Mo, C.H., Wong, M.H., Li, H., 2019. Arbuscular mycorrhizal fungi and the associated bacterial community influence the uptake of cadmium in rice. *Geoderma* 337, 749–757.
- Chhajro, M.A., Fu, Q., Shaaban, M., Rizwan, M.S., Jun, Z., Salam, A., Kubar, K.A., Bashir, S., Hongxing, H., Jamro, G.M., 2018. Identifying the functional groups and the influence of synthetic chelators on Cd availability and microbial biomass carbon in Cd-contaminated soil. *Int. J. Phytoremediat.* 20 (2), 168–174.
- Chien, S.H., Gearhart, M.M., Villagarcía, S., 2011. Comparison of ammonium sulfate with other nitrogen and sulfur fertilizers in increasing crop production and minimizing environmental impact: a review. *Soil Sci.* 176 (7), 327–335.
- Chirakkara, R.A., Cameselle, C., Reddy, K.R., 2016. Assessing the applicability of phytoremediation of soils with mixed organic and heavy metal contaminants. *Rev. Environ. Sci. Biol.* 15 (2), 299–326.
- Chrostowski, P., Durda, J.L., Edelmann, K.G., 1991. The use of natural processes for the control of chromium migration. *Remed. J.* 2 (3), 341–351.
- Chu, W., Chan, K.H., 2003. The mechanism of the surfactant-aided soil washing system for hydrophobic and partial hydrophobic organics. *Sci. Total Environ.* 307 (1–3), 83–92.
- Clemens, S., Palmgren, M.G., Kramer, U., 2002. A long way ahead: understanding and engineering plant metal accumulation. *Trends Plant Sci.* 7 (7), 309–315.
- Coetzee, J.J., Bansal, N., Chirwa, E.M., 2018. Chromium in environment, its toxic effect from chrome-mining and ferrochrome industries, and its possible bioremediation. *Expo. Health* 1–12.
- Conceicao, E.T., Bonotio, D.M., 2006. Radionuclides, heavy metals and fluorine incidence at tapira, phosphate rocks. Brazil and their industrial by-products. *Environ. Pollut.* 139, 232–243.
- Cunningham, S.D., Ow, D.W., 1996. Promises and prospects of phytoremediation. *Plant Physiol.* 110 (3), 715–719.
- Da Conceição Gomes, M.A., Hauser-Davis, R.A., de Souza, A.N., Vitoria, A.P., 2016. Metal

- phytoremediation: general strategies, genetically modified plants and applications in metal nanoparticle contamination. *Ecotoxicol. Environ. Saf.* 134, 133–147.
- Dada, E.O., Njoku, K.I., Osuntoki, A.A., Akinola, M.O., 2015. A review of current techniques of physico-chemical and biological remediation of heavy metals polluted soil. *Ethiop. J. Environ. Stud. Manag.* 8 (5), 606–615.
- Danh, L.T., Truong, P., Mammucari, R., Tran, T., Foster, N., 2009. Vetiver grass, *Vetiveria zizanioides*: a choice plant for phytoremediation of heavy metals and organic wastes. *Int. J. Phytoremediat.* 11 (8), 664–691.
- Davis, C.M., Vincent, J.B., 1997. Chromium oligopeptide activates insulin receptor tyrosine kinase activity. *Biochemistry* 6, 4382–4385.
- De Souza Freitas, E.V., Do Nascimento, C.W.A., 2009. The use of NTA for lead phytoextraction from soil from a battery recycling site. *J. Hazard. Mater.* 171 (1–3), 833–837.
- Dermont, G., Bergeron, M., Mercier, G., Richer-Lafleche, M., 2008. Soil washing for metal removal: a review of physical/chemical technologies and field applications. *J. Hazard. Mater.* 152 (1), 1–31.
- Dhanwal, P., Kumar, A., Dudgeja, S., Chhokar, V., Beniwal, V., 2017. Recent advances in phytoremediation technology. In: Kumar, R., Sharma, A.K., Ahluwalia, S.S. (Eds.), *Advances in Environmental Biotechnology*. Springer, Singapore, pp. 227–241.
- Dhiman, S.S., Selvaraj, C., Li, J., Singh, R., Zhao, X., Kim, D., Kim, J.Y., Kang, Y.C., Lee, J.K., 2016. Phytoremediation of metal-contaminated soils by the hyperaccumulator canola (*Brassica napus* L.) and the use of its biomass for ethanol production. *Fuel* 183, 107–114.
- Dong, R., 2005. Molecular cloning and characterization of a phytochelatin synthase gene, *PvPCS1*, from *Pteris vittata* L. *J. Ind. Microbiol. Biot.* 32, 527–533.
- Edgell, K., 1988. USEPA Method Study 37-SW-846 Method 3050 Acid Diagram of Sediments, Sludges and Soils by AA-Direct Aspiration. EPA/600/4-89/012. U.S. Environmental Protection Agency, Washington, D.C. Retrieved from: [https://cfpub.epa.gov/si/public\\_record\\_Report.cfm?Lab=ORD&dirEntryId=30330](https://cfpub.epa.gov/si/public_record_Report.cfm?Lab=ORD&dirEntryId=30330) (Accessed 25 February 2019).
- Emenike, C.U., Jayanthi, B., Agamuthu, P., Fauziah, S.H., 2018. Biotransformation and removal of heavy metals: a review of phytoremediation and microbial remediation assessment on contaminated soil. *Environ. Res.* 26 (2), 156–168.
- Erakhrumen, A.A., 2007. Phytoremediation: An environmentally sound technology for pollution prevention, control and remediation in developing countries. *Edu. Res. Rev.* 2, 151–156.
- Etesami, H., 2018. Bacterial mediated alleviation of heavy metal stress and decreased accumulation of metals in plant tissues: mechanisms and prospects. *Ecotoxicol. Environ. Saf.* 147, 175–191.
- European Commission, 2001. Disposal and Recycling Routes of Sewage Sludge Part 3. Scientific and Technical Report. European Commission DG Environment.
- European Union, 2002. Heavy Metals in Wastes. European Commission on Environment. <http://ec.europa.eu/environment/waste/studies/pdf/heavymetalsreport.pdf>.
- Evangelou, M.W.H., Ebel, M., Schaeffer, A., 2007. Chelate assisted phytoextraction of heavy metals from soil. Effect, mechanism, toxicity, and fate of chelating agents. *Chemosphere* 68 (6), 989–1003.
- Evanko, F.R., Dzombak, D.A., 1997. Remediation of Metals Contaminated Soils and Groundwater (Tech. Rep. TE-97-01). Groundwater Remediation Technologies Analysis Centre, Pittsburg, Pa, USA.
- Farrell, M., Perkins, W.T., Hobbs, P.J., Griffith, G.W., Jones, D.L., 2010. Migration of heavy metals in soil as influenced by compost amendments. *Environ. Pollut.* 158 (1), 55–64.
- Fawzy, E.M., 2008. Soil remediation using in situ immobilization techniques. *Chem. Ecol.* 24 (2), 147–156.
- Felix, H.R.Z., 1997. Field trials for in situ decontamination of heavy metal polluted soils using crops of metal-accumulating plants. *J. Plant Nutr. Soil Sci.* 160 (4), 525–529.
- Fenyvesi, E., Gruia, K., Morillo, E., Villaverde, J., 2019. Traditional and innovative methods for physical and chemical remediation of soil contaminated with organic contaminants. In: Gruiz, K., Meggyes, T., Fenyvesi, E. (Eds.), *Engineering Tools for Environmental Risk Management: Risk Reduction Technologies and Case Studies*. CRC Press/Balkema, Netherlands, pp. 285–362.
- Flathman, P.E., Lanza, G.R., 1998. Phytoremediation: current views on an emerging green technology. *J. Soil Contam.* 7 (4), 415–432.
- Gao, Y., He, J., Ling, W., Hu, H., Liu, F., 2003. Effects of organic acids on copper and cadmium desorption from contaminated soils. *Environ. Int.* 29 (5), 613–618.
- Garbisu, C., Alkorta, I., 1997. Bioremediation: principles and future. *J. Clean Technol. Environ. Toxicol. Occup. Med.* 6, 351–366.
- Garbisu, C., Alkorta, I., 2001. Phytoextraction: a cost effective plant-based technology for the removal of metals from the environment. *Bioresour. Technol.* 77 (3), 229–236.
- Gautam, P.K., Gautam, R.K., Banerjee, S., Chattopadhyaya, M.C., Pandey, J.D., 2016. Heavy metals in the environment: fate, transport, toxicity and remediation technologies. *Nova Sci. Publ.* 60, 101–130.
- Ghori, N.H., Ghori, T., Hayat, M.Q., Imadi, S.R., Gul, A., Altay, V., Ozturk, M., 2019. Heavy metal stress and responses in plants. *Int. J. Environ. Sci. Technol.* <https://doi.org/10.1007/s13762-019-02215-8>.
- Ghosh, M., Singh, S.P., 2005. A review on phytoremediation of heavy metals and utilization of its byproducts. *Appl. Ecol. Environ. Res.* 3 (1), 1–18.
- Gill, S.S., Anjum, N.A., Hasanuzzaman, M., Gill, R., Trivedi, D.K., Ahmad, I., Pereira, E., Tuteja, N., 2013. Glutathione and glutathione reductase: a boon in disguise for plant abiotic stress defense operations. *Plant Physiol. Biochem.* 70, 204–212.
- Gombert, D., 1994. Soil washing and radioactive contamination. *Environ. Prog.* 13 (2), 138–142.
- GOST, 1983. Environmental Protection, Classification of Chemical Substances for Pollution Control, Standard, Moscow (in Russian). pp. 02–83.
- Grcman, H., Velikonja-Bolta, S., Vodnik, D., Kos, B., Lestan, D., 2001. EDTA enhanced heavy metal phytoextraction: metal accumulation, leaching, and toxicity. *Plant Soil* 235 (1), 105–114.
- Gu, Y.G., Gao, Y.P., Lin, Q., 2016. Contamination, bio-accessibility and human health risk of heavy metals in exposed-lawn soils from 28 urban parks in southern China's largest city, Guangzhou. *Appl. Geochem.* 67, 52–58.
- Gunatilake, S.K., 2015. Methods of removing heavy metals from industrial wastewater. *J. Multidiscip. Eng. Sci. Stud.* 1 (1), 12–18.
- Gupta, P., Diwan, B., 2017. Bacterial exopolysaccharide mediated heavy metal removal: a review on biosynthesis, mechanism and remediation strategies. *Biotechnol. Rep.* 13, 58–71.
- Gupta, N., Khan, D.K., Santra, S.C., 2012. Heavy metal accumulation in vegetables grown in a long-term wastewater-irrigated agricultural land of tropical India. *Environ. Monit. Assess.* 184 (11), 6673–6682.
- Gupta, S.K., Roy, S., Chabukdhara, M., Hussain, J., Kumar, M., 2019. Risk of metal contamination in agriculture crops by reuse of wastewater: an ecological and human health risk perspective. In: Singh, R.P., Kolok, A.S., Bartelt-Hunt, S.L. (Eds.), *Water Conservation, Recycling and Reuse: Issues and Challenges*. Springer, Singapore, pp. 55–79.
- Hadia-e-Fatima, Ahmed, A., 2018. Heavy metal pollution - A mini review. *J. Bacteriol. Mycol.* 6 (3), 179–181. <https://doi.org/10.15406/jbmoa.2018.06.00199>.
- Haroob, B., Ping, A., Pervez, A., Irshad, M., 2018. Characterization of heavy metal in soils as affected by long-term irrigation with industrial wastewater. *J. Water Reuse Desalin.* 9 (1), 47–56.
- Hashimoto, Y., Matsufuru, H., Takaoka, M., Tanida, H., Sato, T., 2009. Impacts of chemical amendment and plant growth on lead speciation and enzyme activities in a shooting range soil: an X-ray absorption fine structure investigation. *J. Environ. Qual.* 38 (4), 1420–1428.
- Hédjij, H., Djebali, W., Belkadhi, A., Cabasson, C., Moing, A., Rolin, D., Brouquisse, R., Gallusci, P., Chaïbi, W., 2015. Impact of long-term cadmium exposure on mineral content of *Solanum lycopersicum* plants: consequences on fruit production. *S. Afr. J. Bot.* 97, 176–181.
- Helmisaari, H.S., Salemaa, M., Derome, J., Kiikkilö, O., Uhlig, C., Nieminen, T.M., 2007. Remediation of heavy metal contaminated forest soil using recycled organic matter and native woody plants. *J. Environ. Qual.* 36 (4), 1145–1153.
- Henry, J.R., 2000. An overview of phytoremediation of lead and mercury, in NNEMS Report. USEPA.
- Herawati, N., Suzuki, S., Hayashi, K., Rivai, I.F., Koyama, H., 2000. Cadmium, copper, and zinc levels in rice and soil of Japan, Indonesia, and China by soil type. *Bull. Environ. Contam. Toxicol.* 64, 33–39.
- Hu, P.J., Qiu, R.L., Senthilkumar, P., Jiang, D., Chen, Z.W., Tang, Y.T., Liu, F.J., 2009. Tolerance, accumulation and distribution of zinc and cadmium in hyperaccumulator *Potentilla griffithii*. *Environ. Exp. Bot.* 66, 317–325.
- Jadia, C.D., Fulekar, H.M., 2008. Phytoremediation: the application of vermicompost to remove zinc, cadmium, copper, nickel and lead by sunflower plant. *Environ. Eng. Manag. J.* 7 (5), 547–558.
- Jaiswal, A., Verma, A., Jaiswal, P., 2018. Detrimental effects of heavy metals in soil, plants, aquatic ecosystem as well as in humans. *J. Environ. Pathol. Toxicol.* 37 (3), 183–197.
- Jamali, M.K., Kazi, T.G., Arain, M.B., Afridi, H.I., Jalbani, N., Memon, A.U.R., Ansari, R., Shah, A., 2007. The feasibility of using an industrial sewage sludge produce in Pakistan as agricultural fertilizer used for cultivation of *Sorghum bicolor* L. *Arch. Agron. Soil Sci.* 53 (6), 659–671.
- Jan, A.T., Azam, M., Siddiqui, K., Ali, A., Choi, I., Haq, Q.M., 2015. Heavy metals and human health: mechanistic insight into toxicity and counter defense system of antioxidants. *Int. J. Mol. Sci.* 16 (12), 29592–29630.
- Jankong, P., Visoottiviset, P., Khokiattiwong, S., 2007. Enhanced phytoremediation of arsenic contaminated land. *Chemosphere* 68, 1906–1912.
- Javed, M.T., Tanwir, K., Akram, M.S., Shahid, M., Niazi, N.K., Lindberg, S., 2019. Phytoremediation of cadmium-polluted water/sediment by aquatic macrophytes: role of plant-induced pH changes. In: Hasanuzzaman, M., Prasad, M.N.V., Fujita, M. (Eds.), *Cadmium Toxicity and Tolerance in Plants*. Academic Press, pp. 495–529.
- Ji, P., Sun, T., Song, Y., Ackland, M.L., Liu, Y., 2011. Strategies for enhancing the phytoremediation of cadmium-contaminated agricultural soils by *Solanum nigrum* L. *Environ. Pollut.* 159, 762–768.
- Jiang, B., Adebayo, A., Jia, J., Xing, Y., Deng, S., Guo, L., Liang, Y., Zhang, D., 2019a. Impacts of heavy metals and soil properties at a Nigerian e-waste site on soil microbial community. *J. Hazard. Mater.* 362, 187–195.
- Jiang, M., Liu, S., Li, Y., Li, X., Luo, Z., Song, H., Chen, Q., 2019b. EDTA-facilitated toxic tolerance, absorption and translocation and phytoremediation of lead by dwarf bamboos. *Ecotoxicol. Environ. Saf.* 170, 502–512.
- Juhasz, A.L., Smith, E., Smith, J., Naidu, R., 2003. In situ remediation of DDT-contaminated soil using a two-phase co-solvent flushing-fungal biosorption process. *Water Air Soil Pollut.* 147 (1–4), 263–274.
- Kalaivanan, D., Ganeshamurthy, A.N., 2016. Mechanisms of heavy metal toxicity in plants. In: Rao, N.K.S., Shivashankara, K.S., Laxman, H.V. (Eds.), *Abiotic Stress Physiology of Horticultural Crops*. Springer, New Delhi, pp. 85–102.
- Kalve, S., Sarangi, B.K., Pandey, R.A., Chakrabarti, T., 2011. Arsenic and chromium hyperaccumulation by an ecotype of *Pteris vittata*-prospective for phytoextraction from contaminated water and soil. *Curr. Sci.* 100, 888–894.
- Kaplan, M., Orman, S., 1998. Effect of elemental sulfur and sulfur containing waste in a calcareous soil in turkey. *J. Plant Nutr.* 21 (8), 1655–1665.
- Karami, A., Shamsuddin, Z.H., 2010. Phytoremediation of heavy metals with several efficiency enhancer methods. *Afr. J. Biotechnol.* 9 (25), 3689–3698.
- Karimzarchi, M., Aminuddin, H., Khanif, M.Y., Radziah, O., 2016. Effect of elemental sulphur timing and application rates on soil P release and concentration in maize. *Pertanika J. Trop. Agric. Sci.* 39 (2), 235–248.
- Kayser, A., Wenger, K., Keller, A., Attinger, W., Felix, H.R., Gupta, S.K., 2000. Enhancement of phytoextraction of Zn, Cd, and Cu from calcareous soil: the use of

- NTA and sulphur amendments. *Environ. Sci. Technol.* 34 (9), 1778–1783.
- Keshavarzi, B., Mokhtarzadeh, Z., Moore, F., RastegariMehri, M., Lahijanzadeh, A., Rostami, S., Kaabi, H., 2015. Heavy metals and polycyclic aromatic hydrocarbons in surface sediments of Karoon River, Khuzestan Province, Iran. *Environ. Sci. Pollut. Res.* 22, 19077–19092.
- Khan, A., Khan, S., Alam, M., Khan, M.A., Aamir, M., Qamar, Z., Rehman, Z.U., Perveen, S., 2016. Toxic metal interactions affect the bioaccumulation and dietary intake of macro- and micro-nutrients. *Chemosphere* 146, 121–128.
- Khalid, S., Shahid, M., Niazi, N.K., Murtaza, B., Bibi, I., Dumat, C., 2017. A comparison of technologies for remediation of heavy metal contaminated soils. *J. Geochem. Explor.* 182, 247–268.
- Kleckerova, A., Docekalova, H., 2014. Dandelion plants as a biomonitor of urban area contamination by heavy metals. *Int. J. Environ. Res.* 8, 157–164.
- Kolacinski, Z., Rincon, J.M., Szymanski, T.P., Sobiecka, E., 2017. Thermal plasma vitrification process as the effective technology for hospital incineration fly ash immobilization. In: Rincon, J.M., Vidal, M.J. (Eds.), *Vitrification and Geopolymerization of Wastes for Immobilization or Recycling*. Universidad Miguel Hernández, pp. 51.
- Kongshaug, G., 1992. Inputs of trace elements to soils and plants. *The Norwegian Academy of Science and Letters, Proc. Chemical Climatology and Geomedical Problems*, Nnrsk Hydro, Oslo, Norway, pp.185–216.
- Kramer, U., 2018. The plants that suck up metal. *Ger. Res.* 40 (3), 18–23.
- Kumar, P.S., Gunasundari, E., 2018. Bioremediation of heavy metals. In: Varjani, S.J., Agarwal, A.K., Gnansounou, E., Gurunathan, B. (Eds.), *Bioremediation: Applications for Environmental Protection and Management*. Springer, Singapore, pp. 165–195.
- Kumar, V., Shahi, S.K., Singh, S., 2018. Bioremediation: an eco-sustainable approach for restoration of contaminated sites. In: Singh, J., Sharma, D., Kumar, G., Sharma, N.R. (Eds.), *Microbial Bioprospecting for Sustainable Development*. Springer, Singapore, pp. 115–136.
- Kumpiene, J., Nordmark, D., Carabante, I., Suziedelyte-Visockiene, J., Aksamitauskas, V.C., 2017. Remediation of soil contaminated with organic and inorganic wood impregnation chemicals by soil washing. *Chemosphere* 184, 13–19.
- Kuppasamy, S., Thavamani, P., Venkateswarlu, K., Lee, Y.B., Naidu, R., Megharaj, M., 2017. Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils: technological constraints, emerging trends and future directions. *Chemosphere* 168, 944–968.
- Kurek, E., Bollag, J.M., 2004. Microbial immobilization of cadmium released from CdO in the soil. *Biogeochem* 69 (2), 227–239.
- Ladislav, S., El-Mufleh, A., Gerente, C., Chazarenc, F., Andres, Y., Bechet, B., 2012. Potential of aquatic macrophytes as bio-indicators of heavy metal pollution in urban storm water runoff. *Water Air Soil Pollut.* 223, 877–888.
- Lajayer, B.A., Moghadam, N.K., Maghsoodi, M.R., Ghorbanpour, M., Kariman, K., 2019. Phytoextraction of heavy metals from contaminated soil, water and atmosphere using ornamental plants: mechanisms and efficiency improvement strategies. *Environ. Sci. Pollut. Res.* <https://doi.org/10.1007/s11356-019-04241-y>.
- Lasat, M.M., 2000. Phytoextraction of metals from contaminated soil: a review of plant/soil/metal interaction and assessment of pertinent agronomic issues. *J. Hazard. Subst. Res.* 2, 1–25.
- Lasat, M.M., 2002. Phytoextraction of toxic metals: a review of biological mechanisms. *J. Environ. Qual.* 31 (1), 109–120.
- Leumann, C.D., Rammelt, R., Gupta, S.K., 1995. Soil remediation by plants: possibilities and limitations. *Sci. Total Environ.* 2, 431–434.
- Lewicki, S., Zdanowski, R., Krzyzowska, M., Lewicka, A., Debski, B., Niemcewicz, M., 2014. The role of chromium III in the organism and its possible use in diabetes and obesity treatment. *Ann. Agric. Environ. Med.* 21, 331–345.
- Leyssens, L., Vinck, B., Van Der Straeten, C., Wuyts, F., Maes, L., 2017. Cobalt toxicity in humans—a review of the potential sources and systemic health effects. *Toxicology* 387, 43–56.
- Li, Y.M., Chaney, R., Brewer, E., Roseberg, R., Angle, J.S., Baker, A., Reeves, R., Nelkin, J., 2003. Development of a technology for commercial phytoextraction of nickel: Economic and technical considerations. *Plant Soil* 249, 107–115.
- Li, Z., Shuman, L.M., 1996. Heavy metal movement in metal contaminated soil profiles. *Soil Sci.* 161, 656–666.
- Li, Y., Zhao, J., Guo, J., Liu, M., Xu, Q., Li, H., Li, Y.F., Zheng, L., Zhang, Z., Gao, Y., 2017. Influence of sulfur on the accumulation of mercury in rice plant (*Oryza sativa* L.) growing in mercury contaminated soils. *Chemosphere* 182, 293–300.
- Liao, X., Li, Y., Yan, X., 2016. Removal of heavy metals and arsenic from a co-contaminated soil by sieving combined with washing process. *J. Environ. Sci.* 41, 202–210.
- Lim, S.L., Lee, L.H., Wu, T.Y., 2016. Sustainability of using composting and vermicomposting technologies for organic solid waste biotransformation: recent overview, greenhouse gases emissions and economic analysis. *J. Clean. Prod.* 111, 262–278.
- Liu, J., Liu, Y.J., Liu, Y., Liu, Z., Zhang, A.N., 2018. Quantitative contributions of the major sources of heavy metals in soils to ecosystem and human health risks: a case study of Yulin, China. *Ecotoxicol. Environ. Saf.* 164, 261–269.
- Liu, W., Zhou, Q., An, J., Sun, Y., Liu, R., 2010. Variations in cadmium accumulation among Chinese cabbage cultivars and screening for Cd-safe cultivars. *J. Hazard. Mater.* 173 (1–3), 737–743.
- Lloyd, J.R., Oremland, R.S., 2006. Microbial transformations of arsenic in the environment: from soda lakes to aquifers. *Elements* 2, 85–90.
- Lockwood, C.L., Stewart, D.I., Mortimer, R.J., Mayes, W.M., Jarvis, A.P., Gruiz, K., Burke, I.T., 2015. Leaching of copper and nickel in soil-water systems contaminated by bauxite residue (red mud) from Ajka, Hungary: the importance of soil organic matter. *Environ. Sci. Pollut. Res.* 22 (14), 10800–10810.
- Long, X.X., Yang, X.E., Ni, W.Z., 2002. Current status and prospective on phytoremediation of heavy metal polluted soils. *J. Appl. Ecol.* 13, 757–762.
- Lu, Y., Luo, D., Lai, A., Liu, G., Liu, L., Long, J., Zhang, H., Chen, Y., 2017. Leaching characteristics of EDTA-enhanced phytoextraction of Cd and Pb by *Zea mays* L. in different particle-size fractions of soil aggregates exposed to artificial rain. *Environ. Sci. Pollut. Res.* 24 (2), 1845–1853.
- Luo, C., Shen, Z., Li, X., 2005. Enhanced phytoextraction of Cu, Pb, Zn and Cd with EDTA and EDDS. *Chemosphere* 59, 1–11.
- Luo, T., Shen, M., Zhou, J., Wang, X., Xia, J., Fu, Z., Jin, Y., 2019. Chronic exposure to low doses of Pb induces hepatotoxicity at the physiological, biochemical, and transcriptomic levels of mice. *Environ. Toxicol.* <https://doi.org/10.1002/tox.22706>.
- Ma, Y., Rajkumar, M., Zhang, C., Freitas, H., 2016. Beneficial role of bacterial endophytes in heavy metal phytoremediation. *J. Environ. Manag.* 174, 14–25.
- Mahar, A., Wang, P., Ali, A., Awasthi, M.K., Lahori, A.H., Wang, Q., Li, R., Zhang, Z., 2016. Challenges and opportunities in the phytoremediation of heavy metals contaminated soils: a review. *Ecotoxicol. Environ. Saf.* 126, 111–121.
- Maimon, A., Khairiah, J., Ahmad, M.R., Aminah, A., Ismail, B.S., 2009. Comparative accumulation of heavy metals in selected vegetables, their availability and correlation in lithogenic and nonlithogenic fractions of soils from some agricultural areas in Malaysia. *Adv. Environ. Biol.* 3, 314–321.
- Malik, N., Biswas, A.K., 2012. Role of higher plants in remediation of metal contaminated sites. *Sci. Rev. Chem. Commun.* 2, 141–146.
- Mandal, P., 2017. An insight of environmental contamination of arsenic on animal health. *Emerg. Contam.* 3 (1), 17–22.
- Mandal, S., Sarkar, B., Bolan, N., Ok, Y.S., Naidu, R., 2017. Enhancement of chromate reduction in soils by surface modified biochar. *J. Environ. Manag.* 186, 277–284.
- Mani, D., Kumar, C., Patel, N.K., 2015. Hyperaccumulator oilcake manure as an alternative for chelate-induced phytoremediation of heavy metals contaminated alluvial soils. *Int. J. Phytoremediat.* 17 (3), 256–263.
- Marrugo-Negrete, J., Pinedo-Hernández, J., Díez, S., 2017. Assessment of heavy metal pollution, spatial distribution and origin in agricultural soils along the Sinú River Basin, Colombia. *Environ. Res.* 154, 380–388.
- Martin, T.A., Ruby, M.V., 2004. Review of in situ remediation technologies for lead, zinc and cadmium in soil. *Remediation* 4 (3), 35–53.
- Martley, E., Gulson, B.L., Pfeifer, H.R., 2004. Metal concentrations in soils around the copper smelter and surrounding industrial complex of Port Kembla, NSW, Australia. *Sci. Total Environ.* 325, 113–127.
- Mattina, M.J.I., Lannucci-Berger, W., Musante, C., White, J.C., 2003. Concurrent plant uptake of heavy metals and persistent organic pollutants from soil. *Environ. Pollut.* 124 (3), 375–378.
- Maturi, M., Reddy, K.R., 2008. Extractants for the removal of mixed contaminants from soils. *Soil Sediment Contam.* 17 (6), 586–608.
- McGrath, S.P., Zhao, F.J., 2003. Phytoextraction of metals and metalloids from contaminated soil. *Curr. Opin. Biotech.* 14, 277–282.
- McGrath, S.P., Zhao, F.J., Lombi, E., 2002. Phytoextraction of metals, metalloids, and radionuclides. *Adv. Agron.* 75, 1–56.
- McLaughlin, M.J., Hamon, R.E., McLaren, R.J., Speir, T.W., Rogers, S.L., 2000. Review: a bioavailability-based rationale for controlling metal and metalloid contamination of agricultural land in Australia and New Zealand. *Aust. J. Soil Res.* 38 (6), 1037–1086.
- McNeil, K.R., Waring, S., 1992. Vitrification of contaminated soils. In: Rees, J.F. (Ed.), *Contaminated Land Treatment Technologies*. Society of Chemical Industry Elsevier, Applied Sciences London, pp. 143–159.
- Meers, E., Ruttens, A., Hopgood, M.J., Samson, D., Tack, F.M.G., 2005. Comparison of EDTA and EDDS as potential soil amendments for enhanced phytoextraction of heavy metals. *Chemosphere* 58, 1011–1022.
- Mesjasz-Przybyłowicz, J., Nakonieczny, M., Migula, P., Augustyniak, M., Tarnawska, M., Reimold, W.U., Koerberl, C., Przybyłowicz, W., Glowacka, E., 2004. Uptake of cadmium, lead, nickel and zinc from soil and water solutions by the nickel hyperaccumulator *Berkeya coddii*. *Acta Biol. Cracov. Ser. Bot.* 46, 75–85.
- Missimer, T., Teaf, C., Beeson, W., Maliva, R., Wooschlager, J., Covert, D., 2018. Natural background and anthropogenic arsenic enrichment in florida soils, surface water, and groundwater: a review with a discussion on public health risk. *Int. J. Environ. Res. Public Health* 15 (10), 1–30.
- Monni, S., Salemaa, M., Miller, N., 2000. The Tolerance of *Empetrum nigrum* to Copper and Nickel. *Environ. Pollut.* 109, 221–229.
- Morcillo, P., Esteban, M.A., Cuesta, A., 2016. Heavy metals produce toxicity, oxidative stress and apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere* 144, 225–233.
- Mortensen, L.H., Rønn, R., Vestergård, M., 2018. Bioaccumulation of cadmium in soil organisms—With focus on wood ash application. *Ecotoxicol. Environ. Saf.* 156, 452–462.
- Mukesh, K.R., Kumar, P., Singh, M., Singh, A., 2008. Toxic effect of heavy metals in livestock health. *Veterin. World* 1, 28–30.
- Mukhopadhyay, S., Maiti, S.K., 2010. Phytoremediation of metal enriched mine waste: a review. *Glob. J. Environ. Res.* 4, 135–150.
- Mukhopadhyay, R., Rosen, B.P., Phung, L.T., Silver, S., 2002. Microbial arsenic: from geocycles to genes and enzymes. *FEMS Microbiol. Rev.* 26, 311–325.
- Murtaza, G., Javed, W., Hussain, A., Wahid, A., Murtaza, B., Owens, G., 2015. Metal uptake via phosphate fertilizer and city sewage in cereal and legume crops in Pakistan. *Environ. Sci. Pollut. Res.* 22 (12), 9136–9147.
- Mwegha, W.J.S., 2008. The use of phytoremediation technology for abatement soil and groundwater pollution in Tanzania: opportunities and challenges. *J. Sustain. Dev. Afr.* 10 (1), 140–156.
- Naderizadeh, Z., Khademi, H., Ayoubi, S., 2016. Biomonitoring of atmospheric heavy metals pollution using dust deposited on date palm leaves in southwestern Iran. *Atmosfera* 29 (2), 141–155.
- Naeem, M., Ali, Q., Shahbaz, M., Ali, F., 2011. Role of rhizobacteria in phytoremediation of heavy metals: an overview. *Int. Res. J. Plant Sci.* 2, 220–232.
- Nazir, R., Khan, M., Masab, M., Rehman, H.U., Rauf, N.U., Shahab, S., Ameer, N., Sajed, M.,

- Ullah, M., Rafeeq, M., Shaheen, Z., 2015. Accumulation of heavy metals (Ni, Cu, Cd, Cr, Pb, Zn, Fe) in the soil, water and plants and analysis of physico-chemical parameters of soil and water collected from Tanda Dam Kohat. *J. Pharm. Sci. Res.* 7 (3), 89–97.
- Nealson, K.H., Rosson, R.A., Mayers, C.R., 1992. Microbial reduction of manganese and iron: new approaches to carbon cycling. *Appl. Environ. Microb.* 58, 439–443.
- Nies, D.H., 2003. Efflux-mediated heavy metal resistance in prokaryotes. *FEMS Microbiol. Rev.* 27, 313–339.
- Nie, S.W., Gao, W.S., Chen, Y.Q., Sui, P., Eneji, A.E., 2010. Use of life cycle assessment methodology for determining phytoremediation potentials of maize-based cropping systems in fields with nitrogen fertilizer over-dose. *J. Clean. Prod.* 18 (15), 1530–1534.
- Nishimune, A., 1993. Combinations of organic and inorganic fertilizers for crop production. *Ext. Bull.* 372, 1–25.
- Odoemelam, S.A., Ukpe, R.A., 2008. Heavy metal decontamination of polluted soils using *Bryophyllum pinnatum*. *Afr. J. Biotechnol.* 7 (23), 4301–4303.
- Ogunleye, P.O., Mayaki, M.C., Amapu, I.Y., 2002. Radioactivity and heavy metal composition of Nigerian phosphate rocks. Possible environmental implications. *J. Environ. Radioact.* 62, 39–48.
- Oves, M., Khan, M.S., Qari, A.H., Felemban, M.N., Almeelbi, T., 2016. Heavy metals: biological importance and detoxification strategies. *J. Bioremed. Biodeg.* 7 (2), 1–15.
- Padmavathiamma, P.K., Li, L.Y., 2007. Phytoremediation technology: hyperaccumulation metals in plants. *Water Air Soil Pollut.* 184, 105–126.
- Pajević, S., Borisav, M., Nikolic, N., Arsenov, D.D., Orlovic, S., Zupunski, M., 2016. Phytoextraction of heavy metals by fast-growing trees: a review. In: Tsao, D. (Ed.), *Phytoremediation*. Springer-Verlag Berlin Heidelberg, pp. 29–64.
- Pandey, P., Dubey, R.S., 2019. Metal toxicity in rice and strategies for improving stress tolerance. In: Hasanuzzaman, M., Fujita, M., Nahar, K., Biswas, J.K. (Eds.), *Advances in Rice Research for Abiotic Stress Tolerance*. Woodhead Publishing, pp. 313–339.
- Pantelica, A.L., Salagean, M.N., 1997. INAA of some phosphates used in fertilizer industries. *J. Radioanal. Nucl. Chem.* 216, 261–264.
- Park, J., Jung, Y., Han, M., Lee, S., 2002. Simultaneous removal of cadmium and turbidity in contaminated soil-washing water by DAF and electroflotation. *Water Sci. Technol.* 46 (11–12), 225–230.
- Parmar, S., Singh, V., 2015. Phytoremediation approaches for heavy metal pollution: a review. *J. Plant Sci. Res.* 2 (2), 139.
- Patra, D.K., Pradhan, C., Patra, H.K., 2018. Chelate based phytoremediation study for attenuation of chromium toxicity stress using lemongrass: *Cymbopogon flexuosus* (nees ex steud.) W. Watson. *Int. J. Phytoremediat.* 20 (13), 1324–1329.
- Paulo, L.M., Ramiro-Garcia, J., van Mourik, S., Stams, A.J., Sousa, D.Z., 2017. Effect of nickel and cobalt on methanogenic enrichment cultures and role of biogenic sulfide in metal toxicity attenuation. *Front. Microbiol.* 8, 1–12.
- Pence, N.S., Larsen, P.B., Ebbs, S.D., Latham, D.L., Lasat, M.M., Garvin, D.F., Eide, D., Kochian, L.V., 2000. The molecular physiology of heavy metal transport in the Zn/Cd hyperaccumulator *Thlaspi caerulescens*. *Proc. Natl. Acad. Sci. USA* 97, 4956–4960.
- Perrier, F., Yan, B., Candaudap, F., Pokrovsky, O.S., Gourdain, E., Meleard, B., Bussiere, S., Coriou, C., Robert, T., Nguyen, C., Cornu, J.Y., 2016. Variability in grain cadmium concentration among durum wheat cultivars: impact of aboveground biomass partitioning. *Plant Soil* 404 (1–2), 307–320.
- Pilon-Smits, E., 2005. Phytoremediation. *Annu. Rev. Plant Biol.* 56, 15–39.
- Postigo, C., Martinez, D.E., Grondona, S., Miglioranza, K.S.B., 2017. Groundwater Pollution: Sources, Mechanisms, and Prevention. Reference Module in Earth Systems and Environmental Sciences. Elsevier, New York, pp. 143–162.
- Prieto, M.J., Acevedo, S.O.A., Prieto, G.F., 2018. Phytoremediation of soils contaminated with heavy metals. *Biodivers. Int. J.* 2 (4), 362–376. <https://doi.org/10.15406/bij.2018.02.00088>.
- Qian, S.Q., Liu, Z., 2000. An overview of development in the soil-remediation technologies. *Chem. Ind. Eng. Pro.* 4, 10–12.
- Qiao, J., Sun, H., Luo, X., Zhang, W., Mathews, S., Yin, X., 2017. EDTA-assisted leaching of Pb and Cd from contaminated soil. *Chemosphere* 167, 422–428.
- Qureshi, A.S., Hussain, M.I., Ismail, S., Khan, Q.M., 2016. Evaluating heavy metal accumulation and potential health risks in vegetables irrigated with treated wastewater. *Chemosphere* 163, 54–61.
- RAAG, 2000. Evaluation of Risk Based Corrective Action Model Remediation. Alternative Assessment Group, Memorial University of Newfoundland, St John's, NF, Canada.
- Rai, P.K., 2008. Phytoremediation of Hg and Cd from industrial effluents using an aquatic free floating macrophyte *Azolla pinnata*. *Int. J. Phytorem.* 10, 430–439.
- Rai, R., Agrawal, M., Agrawal, S.B., 2016. Impact of heavy metals on physiological processes of plants: with special reference to photosynthetic system. In: Singh, A., Prasad, S.M., Singh, R.P. (Eds.), *Plant Responses to Xenobiotics*. Springer, Singapore, pp. 127–140.
- Rakshae, R., Giah, M., Pourahmad, A., 2009. Studying effect of cell wall's carboxyl-carboxylate ratio change of Lemna minor to remove heavy metals from aqueous solution. *J. Hazard. Mater.* 163 (1), 165–173.
- Ramamurthy, A.S., Memarian, R., 2012. Phytoremediation of mixed soil contaminants. *Water Air Soil Pollut.* 223, 511–518.
- Ramanjaneyulu, A.V., Neelima, T.L., Madhavi, A., Ramprakash, T., 2017. Phytoremediation: an overview. In: Humberto, R.M., Ashok, G.R., Thakur, K., Sarkar, N.C. (Eds.), *Applied Botany*. American Academic Press, pp. 42–84.
- Raskin, I., Ensley, B.D., 2000. *Phytoremediation of Toxic Metals: Using Plants to Clean up the Environment*, first ed. John Wiley and Sons, Inc, New York.
- Reed, S.C., Crites, R.W., Middlebrooks, E.J., 1995. *Natural Systems for Waste Management and Treatment*, second ed. McGraw-Hill, New York, NY, USA.
- Reeves, R.D., Baker, A.J.M., 2000. Metal-accumulating plants. In: Raskin, I., Ensley, B.D. (Eds.), *Phytoremediation of Toxic Metals: Using Plants to Clean-Up the Environment*. John Wiley and Sons, New York, pp. 193–230.
- Riser-Roberts, E., 1998. Remediation of Petroleum Contaminated Soil: Biological, Physical, and Chemical Processes, first ed. Lewis Publishers, Boca Raton, FL.
- Rodriguez, L., Lopez-Bellido, F.J., Carnicer, A., Recreo, F., Tallos, A., Monteagudo, J.M., 2005. Mercury recovery from soils by phytoremediation. In: *Book of Environmental Chemistry*. Springer, Berlin, Germany, pp. 197–204.
- Romkens, P., Bouwman, L., Japenga, J., Draaisma, C., 2002. Potentials and drawbacks of chelate-enhanced phytoremediation of soils. *Environ. Pollut.* 116, 109–121.
- Rosen, C.J., 2002. Lead in the Home Garden and Urban Soil Environment. Communication and Educational Technology Services, University of Minnesota Extension.
- RoyChowdhury, A., Datta, R., Sarkar, D., 2018. Heavy metal pollution and remediation. In: Torok, B., Dransfield, T. (Eds.), *Green Chemistry*. Elsevier, pp. 359–373.
- Ryzhenko, N.O., Kavetsky, S.V., Kavetsky, V.M., 2017. Cd, Zn, Cu, Pb, Co, Ni phytotoxicity assessment. *Pol. J. Soil Sci.* 50 (2), 197–215.
- Sabiha, J., Mehmood, T., Chaudhry, M.M., Tufail, Irfan, N., 2009. Heavy metal pollution from phosphate rock used for the production of fertilizer in Pakistan. *Microchem. J.* 91, 94–99.
- Saha, M., Sarkar, S., Sarkar, B., Sharma, B.K., Bhattacharjee, S., Tribedi, P., 2016. Microbial siderophores and their potential applications: a review. *Environ. Sci. Pollut. Res.* 23 (5), 3984–3999.
- Saif, S., Khan, M.S., 2017. Assessment of heavy metals toxicity on plant growth promoting rhizobacteria and seedling characteristics of *Pseudomonas putida* SFB3 inoculated greengram. *Acta Sci. Agric.* 1, 47–56.
- Saifullah, Meers, E., Qadir, M., De-Caritat, P., Tack, F.M.G., Du-Laing, G., Zia, M.H., 2009. EDTA-assisted Pb phytoextraction. *Chemosphere* 74 (10), 1279–1291.
- Sakakibara, M., Ohmori, Y., Ha, N.T.H., Sano, S., Sera, K., 2011. Phytoremediation of heavy metal contaminated water and sediment by *Eleocharis acicularis*. *Clean Soil Air Water* 39, 735–741.
- Sakizadeh, M., Ghorbani, H., 2017. Concentration of heavy metals in soil and staple crops and the associated health risk. *Arch. Hyg. Sci.* 6 (4), 303–313.
- Sandeep, G., Vijayalatha, K.R., Anitha, T., 2019. Heavy metals and its impact in vegetable crops. *Int. J. Chem. Stud.* 7 (1), 1612–1621.
- Sarma, H., 2011. Metal hyperaccumulation in plants: a review focusing on phytoremediation technology. *Environ. Sci. Technol.* 4, 118–138.
- Sarubbo, L.A., Rocha Jr, R.B., Luna, J.M., Rufino, R.D., Santos, V.A., Banat, I.M., 2015. Some aspects of heavy metals contamination remediation and role of biosurfactants. *Chem. Ecol.* 31 (8), 707–723.
- Sarwar, N., Imran, M., Shaheen, M.R., Ishaque, W., Kamran, M.A., Matloob, A., Rehman, A., Hussain, S., 2017. Phytoremediation strategies for soils contaminated with heavy metals: modifications and future perspectives. *Chemosphere* 171, 710–721.
- Schalk, I.J., Hannauer, M., Braud, A., 2011. New roles for bacterial siderophores in metal transport and tolerance. *Environ. Microbiol.* 13, 2844–2854.
- Schnoor, J.L., 1997. *Phytoremediation*. University of Iowa, Department of Civil and Engineering, pp. 1–62.
- Schnoor, J.L., 2000. Phytostabilization of metals using hybrid poplar trees. In: Raskin, I., Ensley, B.D. (Eds.), *Phytoremediation of Toxic Metals: Using Plants to Clean-up the Environment*. John Wiley and Sons, New York, pp. 133–150.
- Schutzenbeutel, A., Polle, A., 2002. Plant responses to abiotic stresses: heavy metal-induced oxidative stress and protection by mycorrhization. *J. Exp. Bot.* 53, 1351–1365.
- Schwartz, C., Echevarria, G., Morel, J.L., 2003. Phytoextraction of cadmium with *Thlaspi caerulescens*. *Plant Soil* 249 (1), 27–35.
- Schwartz, G.E., Rivera, N., Lee, S.W., Harrington, J.M., Hower, J.C., Levine, K.E., Vengosh, A., Hsu-Kim, H., 2016. Leaching potential and redox transformations of arsenic and selenium in sediment microcosms with fly ash. *Appl. Geochem.* 67, 177–185.
- Scragg, A., 2006. *Environmental Biotechnology*, second ed. Oxford University Press, UK.
- Seidel, H., Ondruschka, J., Morgenstern, P., Stottmeister, U., 1998. Bioleaching of heavy metals from contaminated aquatic sediments using indigenous sulphur oxidizing bacteria: a feasibility study. *Water Sci. Technol.* 37, 387–394.
- Sheoran, V., Sheoran, A., Poonia, P., 2011. Role of hyperaccumulators in phytoremediation of metals from contaminated mining sites: a review. *Crit. Rev. Environ. Sci. Technol.* 41, 168–214.
- Sidhu, G.P.S., 2016. Heavy metal toxicity in soils: sources, remediation technologies and challenges. *Adv. Plants Agric. Res.* 5 (1), 1–3.
- Sidhu, G.P.S., Bali, A.S., Singh, H.P., Batish, D.R., Kohli, R.K., 2018. Ethylenediamine disuccinic acid enhanced phytoextraction of nickel from contaminated soils using *Coronopus didymus* (L.) Sm. *Chemosphere* 205, 234–243.
- Singh, S., 2012. Phytoremediation: a sustainable alternative for environmental challenges. *Int. J. Gr. Herb. Chem.* 1, 133–139.
- Singh, V., Bhargava, M., 2017a. Phytomining: principles and Applications. In: Bhargava, A., Srivastava, S. (Eds.), *Biotechnology: Recent Trends and Emerging Dimensions*. CRC Press, pp. 141–159.
- Singh, V., Bhargava, M., 2017b. Phytomining: principles and Applications. In: Bhargava, A., Srivastava, S. (Eds.), *Biotechnology: Recent Trends and Emerging Dimensions*. CRC Press, pp. 141–159.
- Singh Brar, B., Singh, J., Singh, G., Kaur, G., 2015. Effects of long-term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize-wheat rotation. *Agron* 5 (2), 220–238.
- Smith, L.A., Means, J.L., Chen, A., 1995. *Remedial Options for Metals-Contaminated Sites*. Lewis Publishers, Boca Raton, Fla, USA.
- Smolen, S., Sady, W., Ledwozyw-Smolen, I., 2010. Quantitative relations between the content of selected trace elements in soil extracted with 0.03M CH<sub>3</sub>COOH or 1M HCl and its total concentration in lettuce and spinach. *Acta Sci. Pol. Hortorum Cultus* 4, 13–23.
- Smolinska, B., 2015. Green waste compost as an amendment during induced phytoextraction of mercury-contaminated soil. *Environ. Sci. Pollut. Res.* 22 (5), 3528–3537.
- Song, Y., Amami, M.T., Benamar, A., Mezazigh, S., Wang, H., 2016. Effect of EDTA,

- EDDS, NTA and citric acid on electrokinetic remediation of As, Cd, Cr, Cu, Ni, Pb and Zn contaminated dredged marine sediment. *Environ. Sci. Pollut. Res* 23 (11), 10577–10586.
- Song, B., Zeng, G., Gong, J., Liang, J., Xu, P., Liu, Z., Zhang, Y., Zhang, C., Cheng, M., Liu, Y., Ye, S., 2017. Evaluation methods for assessing effectiveness of in situ remediation of soil and sediment contaminated with organic pollutants and heavy metals. *Environ. Int.* 105, 43–55.
- Srivastava, M., Ma, L.Q., Santos, J.A.G., 2006. Three new arsenic hyperaccumulating ferns. *Sci. Total Environ.* 364, 24–31.
- Srivastava, V., Sarkar, A., Singh, S., Singh, P., de Araujo, A.S., Singh, R.P., 2017. Agroecological responses of heavy metal pollution with special emphasis on soil health and plant performances. *Front. Environ. Sci.* 5, 64.
- Sun, G., Sommar, J., Feng, X., Lin, C.J., Ge, M., Wang, W., Yin, R., Fu, X., Shang, L., 2016. Mass-dependent and-independent fractionation of mercury isotope during gas-phase oxidation of elemental mercury vapor by atomic Cl and Br. *Environ. Sci. Technol.* 50 (17), 9232–9241.
- Sun, L., Xue, Y., Peng, C., Xu, C., Shi, J., 2018. Does sulfur fertilizer influence Cu migration and transformation in colloids of soil pore water from the rice (*Oryza sativa* L.) rhizosphere? *Environ. Pollut.* 243, 1119–1125.
- Suthersan, S.S., 1997. Remediation Engineering: Design Concepts. Lewis Publishers, Boca Raton, FL, USA, pp. 27–70.
- Suthersan, S.S., Horst, J., Schnobrich, M., Welty, N., Mc Donough, J., 2016. Remediation Engineering: Design Concepts. CRC Press.
- Suvarapu, L.N., Baek, S.O., 2017. Determination of heavy metals in the ambient atmosphere: a review. *Toxicol. Ind. Health* 33 (1), 79–96.
- Sytar, O., Brestic, M., Taran, N., Zivcak, M., 2016. Plants used for biomonitoring and phytoremediation of trace elements in soil and water. In: Ahmad, P. (Ed.), *Plant Metal Interaction*. Elsevier, pp. 361–384.
- Szynkowska, M.I., Pawlaczyk, A., MacKiewicz, E., 2018. Bioaccumulation and biomagnification of trace elements in the environment. In: Chojnacka, K., Saeid, A. (Eds.), *Recent Advances in Trace Elements*. John Wiley & Sons Ltd, New Jersey, pp. 251–276.
- Tahmasbian, I., Sinegani, A.A.S., 2016. Improving the efficiency of phytoremediation using electrically charged plant and chelating agents. *Environ. Sci. Pollut. Res* 23 (3), 2479–2486.
- Tang, X., Li, X., Liu, X., Hashmi, M.Z., Xu, J., Brookes, P.C., 2015. Effects of inorganic and organic amendments on the uptake of lead and trace elements by *Brassica chinensis* grown in an acidic red soil. *Chemosphere* 119, 177–183.
- Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K., Sutton, D.J., 2012. Heavy metal toxicity and the environment. In: Luch, A. (Ed.), *Molecular, Clinical and Environmental Toxicology*. Basel, Springer, pp. 133–164.
- Tian, Y., Zhang, H., 2016. Producing biogas from agricultural residues generated during phytoremediation process: possibility, threshold, and challenges. *Int. J. Green Energy* 13 (15), 1556–1563.
- Tichy, R., Fajtl, J., Kuzel, S., Kolar, L., 1997. Use of elemental sulphur to enhance a cadmium solubilization and its vegetative removal from contaminated soil. *Nutr. Cycl. Agroecosyst.* 46, 249–255.
- Tripathi, V., Edrisi, S.A., Abhilash, P.C., 2016. Towards the coupling of phytoremediation with bioenergy production. *Renew. Sustain. Energy Rev.* 57, 1386–1389.
- Tripathi, D.K., Singh, S., Singh, S., Mishra, S., Chauhan, D.K., Dubey, N.K., 2015. Micronutrients and their diverse role in agricultural crops: advances and future prospective. *Acta Physiol. Plant.* 37 (7), 1–14.
- Tsai, L.J., Yu, K.C., Chen, S.F., Kung, P.Y., Chang, C.Y., Lin, C.H.P., 2003. Rationing of heavy metals in contaminated river sediment via bioleaching: effect of sulfur added to total solid ratio. *Water Res.* 37, 4623–4630.
- ul-Hassan, Z., Ali, S., Rizwan, M., Hussain, A., Akbar, Z., Rasool, N., Abbas, F., 2017. Role of zinc in alleviating heavy metal stress. In: Naem, M., Ansari, A.A., Gill, S.S. (Eds.), *Essential Plant Nutrients: Uptake, Use Efficiency, and Management*. Springer International Publishing, Switzerland, pp. 351–366.
- United States Environmental Protection Agency (US EPA), 2016. Health Effects Assessment Summary Tables (HEAST); Annual Update. United States Environmental Protection Agency, Washington, DC, USA (2016).
- Urrialdel, V., Albuquerque, B., Guirao-Abad, J.P., Pla, J., Argüelles, J.C., Alonso-Monge, R., 2017. Arsenic inorganic compounds cause oxidative stress mediated by the transcription factor PHO4 in *Candida albicans*. *Microbiol. Res.* 203, 10–18.
- Urum, K., Pekdemir, T., Gopur, M., 2003. Optimum conditions for washing of crude oil-contaminated soil with biosurfactant solutions. *Process Saf. Environ. Prot.* 81 (3), 203–209.
- USEPA, 1992. Vitrification Technologies for Treatment of Hazardous and Radioactive Waste Handbook, Tech. Rep. EPA/625/R-92/002. United States Environmental Protection Agency, Office of Research and Development, Washington, D.C., USA.
- Usman, K., Al-Ghouthi, M.A., Abu-Dieyeh, M.H., 2018. Phytoremediation: halophytes as promising heavy metal hyperaccumulators. In: Saleh, H.E.M., Aqlan, R. (Eds.), *Heavy Metals*. IntechOpen, pp. 201–217.
- Van Gestel, C.A.M., Dirven-Van Breemen, E.M., Kamerman, J.W., 1992. Evaluation of Decontaminated Soils. National Institute of Public Health and Environmental Protection, Bilthoven.
- Van Nevel, L., Mertens, J., Oorts, K., Verheyen, K., 2007. Phytoextraction of metals from soils: how far from practice? *Environ. Pollut.* 150 (1), 34–40.
- Vandenove, H., Van Hees, M., Van Winkel, S., 2001. Feasibility of phytoextraction to clean up low level uranium contaminated soil. *Int. J. Phytoremediat.* 3, 301–320.
- Vangronsveld, J., Herzig, R., Weyens, N., Boulet, J., Adriaensen, J., Ruttens, A., Thewys, T., Vassilev, A., Meers, E., Nehnevajova, E., Van der Lelie, D., Mench, M., 2009. Phytoremediation of contaminated soils and groundwater: lessons from the field. *Environ. Sci. Pollut. Res.* 16, 765–794.
- Vassilev, A., Schwitzgebel, J.P., Thewys, T., Van Der Lelie, D., Vangronsveld, J., 2004. The use of plants for remediation of metal-contaminated soils. *Sci. World J.* 4, 9–34.
- Vishnoi, S.R., Srivastava, P.N., 2008. Phytoremediation-green for environmental clean. In: *Proceedings of the 12th World Lake Conference*, Jaipur, India, pp. 1016–1021.
- Volland, S., Bayer, E., Baumgartner, V., Andosch, A., Lütz, C., Sima, E., Lütz-Meindl, U., 2014. Rescue of heavy metal effects on cell physiology of the algal model system *Micrasterias* by divalent ions. *J. Plant Physiol.* 171 (2), 154–163.
- Wait, S.T., Thomas, D., 2003. The characterization of base oil recovered from the low temperature thermal desorption of drill cuttings, SPE/EPA exploration and production. In: *Environmental Conference*, 10–12 Mar. San Antonio, TX, pp. 151–158.
- Wallace, A., Wallace, G.A., 1993. A possible flaw in EPA, new sludge rule, due to heavy metal interactions. *Commun. Soil Sci. Plant Anal.* 25, 129–135.
- Wan, X., Lei, M., Chen, T., 2016. Cost-benefit calculation of phytoremediation technology for heavy-metal-contaminated soil. *Sci. Total Environ.* 563, 796–802.
- Wang, H., Hu, Y.F., Hao, J.H., Chen, Y.H., Su, P.Y., Wang, Y., Yu, Z., Fu, L., Xu, Y.Y., Zhang, C., Tao, F.B., 2015. Maternal zinc deficiency during pregnancy elevates the risks of fetal growth restriction: a population-based birth cohort study. *Sci. Rep.* 5, 1–10.
- Watanabe, T., Hirano, S., 2013. Metabolism of arsenic and its toxicological relevance. *Arch. Toxicol.* 87 (6), 969–979.
- Watanabe, M.E., 1997. Phytoremediation on the brink of commercialization. *Environ. Sci. Technol.* 31, 182–186.
- Wei, Q., Li, Y., Zu, Y.Q., 2008a. Advances in researching cultivation techniques on hyperaccumulators phytoremediating the heavy metal contaminated soil. *J. Yunnan Agric. Univ.* 23 (1), 104–108.
- Wei, S.H., Teixeira da Silva, J.A., Zhou, Q.X., 2008b. Agro-improving method of phytoextracting heavy metal contaminated soil. *J. Hazard. Mater.* 150, 662–668.
- Wei, B., Yang, L., 2010. A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchem. J.* 94 (2), 99–107.
- Wei, S.H., Zhou, Q.X., 2004. Discussion on basic principles and strengthening measures for phytoremediation of soils contaminated by heavy metals. *Chin. J. Ecol.* 23 (1), 65–72.
- Wenger, K., Gupta, S.K., Furrer, G., Schulin, R., 2003. The role of nitrilotriacetate in copper uptake by tobacco. *J. Environ. Qual.* 32, 1669–1676.
- Wenzel, W.W., Unterbrunner, R., Sommer, P., Pasqualina, S., 2003. Chelate-assisted phytoextraction using canola (*Brassica napus* L.) in outdoors pot and lysimeter experiments. *Plant Soil* 249, 83–96.
- WHO/FAO, 2007. Joint FAO/WHO Food Standard Programme Codex Alimentarius Commission 13th Session. Report of the Thirty Eight Session of the Codex Committee on Food Hygiene. Houston, United States of America, ALINORM 07/30/13.
- Wijayawardena, M.A.A., Megharaj, M., Naidu, R., 2016. Exposure, toxicity, health impacts, and bioavailability of heavy metal mixtures. In: Donald, L.S. (Ed.), *Advances in Agronomy* (V-138). Elsevier Inc, Amsterdam, Netherlands, pp. 175–234.
- Wijayawardena, M.A., Naidu, R., Megharaj, M., Lamb, D., Thavamani, P., Kuchel, T., 2015. Using soil properties to predict in vivo bioavailability of lead in soils. *Chemosphere* 138, 422–428.
- Wiszniewska, A., Hanus-Fajerska, E., Muszynska, E., Ciarkowska, K., 2016. Natural organic amendments for improved phytoremediation of polluted soils: a review of recent progress. *Pedosphere* 26 (1), 1–12.
- Woodford, Chris, 2019. Land pollution. Retrieved from <<https://www.explainthatstuff.com/land-pollution.html>>. (Accessed 29 January 2019).
- Wu, Z., Liu, S., Zhao, J., Wang, F., Du, Y., Zou, S., Li, H., Wen, D., Huang, Y., 2017. Comparative responses to silicon and selenium in relation to antioxidant enzyme system and the glutathione-ascorbate cycle in flowering Chinese cabbage (*Brassica campestris* L. ssp. *chinensis* var. *utilis*) under cadmium stress. *Environ. Exp. Bot.* 133, 1–11.
- Wu, H., Tang, S., Zhang, X., Guo, J., Song, A., Tian, S., Smith, D.L., 2009. Using elevated CO<sub>2</sub> to increase the biomass of a *Sorghum vulgare* x *Sorghum vulgare* var. Sudanense hybrid and *Trifolium pratense* L. and to trigger hyperaccumulation of cesium. *J. Hazard. Mater.* 170, 861–870.
- Wu, Q., Wang, S., Thangavel, P., Li, Q., Zheng, H., Bai, J., Qiu, R., 2011. Phytostabilization potential of *Jatropha curcas* L. in polymetallic acid mine tailings. *Int. J. Phytorem.* 13, 788–804.
- Wuana, R.A., Okieimen, F.E., 2011. Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. *ISRN Ecol.* 2011, 1–20.
- Xia, X., Lin, S., Zhao, J., Zhang, W., Lin, K., Lu, Q., Zhou, B., 2018a. Toxic responses of microorganisms to nickel exposure in farmland soil in the presence of earthworm (*Eisenia fetida*). *Chemosphere* 192, 43–50.
- Xia, W.Y., Feng, Y.S., Du, Y.J., Reddy, K.R., Wei, M.L., 2018b. Solidification and stabilization of heavy metal-contaminated industrial site soil using KMP binder. *J. Mater. Civ. Eng.* 30 (6), 1–9.
- Xia, W.Y., Feng, Y.S., Jin, F., Zhang, L.M., Du, Y.J., 2017. Stabilization and solidification of a heavy metal contaminated site soil using a hydroxyapatite-based binder. *Constr. Build. Mater.* 156, 199–207.
- Xia, S., Song, Z., Jeyakumar, P., Shaheen, S.M., Rinklebe, J., Ok, Y.S., Bolan, N., Wang, H., 2019. A critical review on bioremediation technologies for Cr (VI)-contaminated soils and wastewater. *Crit. Rev. Environ. Sci. Technol.* 1–52.
- Xu, Q., Shi, G., 2000. The toxic effects of single Cd and interaction of Cd with Zn on some physiological index of *Oenanthe javanica*. *J. Nanjing Univ.* 23 (4), 97–100.
- Yadav, R., Arora, P., Kumar, S., Chaudhury, A., 2010. Perspectives for genetic engineering of poplars for enhanced phytoremediation abilities. *Ecotoxicology* 19, 1574–1588.
- Yahaghi, Z., Shirvani, M., Nourbakhsh, F., De La Pena, T.C., Pueyo, J.J., Talebi, M., 2018. Isolation and characterization of pb-solubilizing bacteria and their effects on pb uptake by *Brassica juncea*: implications for microbe-assisted phytoremediation. *J. Microbiol. Biotechnol.* 28 (7), 1156–1167.
- Yan, X., Liu, M., Zhong, J., Guo, J., Wu, W., 2018. How human activities affect heavy

- metal contamination of soil and sediment in a long-term reclaimed area of the Liaohe River Delta, North China. *Sustainable* 10 (2), 1–19.
- Yanai, J., Zhao, F.J., McGrath, S.P., Kosaki, T., 2006. Effect of soil characteristics on Cd uptake by the hyperaccumulator *Thlaspi caerulescens*. *Environ. Pollut.* 139 (1), 167–175.
- Yang, S.X., Deng, H., Li, M.S., 2008. Manganese uptake and accumulation in a woody hyperaccumulator, *Schima superba*. *Plant Soil Environ.* 54, 441–446.
- Yang, Z., Shi, W., Yang, W., Liang, L., Yao, W., Chai, L., Gao, S., Liao, Q., 2018. Combination of bioleaching by gross bacterial biosurfactants and flocculation: a potential remediation for the heavy metal contaminated soils. *Chemosphere* 206, 83–91.
- Yang, Q., Wang, L., Ma, H., Yu, K., Martin, J.D., 2016a. Hydrochemical characterization and pollution sources identification of groundwater in Salawusu aquifer system of Ordos Basin, China. *Environ. Pollut.* 216, 340–349.
- Yang, Z., Zhang, Z., Chai, L., Wang, Y., Liu, Y., Xiao, R., 2016b. Bioleaching remediation of heavy metal-contaminated soils using *Burkholderia* sp. Z-90. *J. Hazard. Mater.* 301, 145–152.
- Ye, R., Wright, A.L., Orem, W.H., McCray, J.M., 2010. Sulfur distribution and transformations in Everglades Agricultural Area soil as influenced by sulfur amendment. *Soil Sci.* 175 (6), 263–269.
- Zarschler, K., Rocks, L., Licciardello, N., Boselli, L., Polo, E., Garcia, K.P., De Cola, L., Stephan, H., Dawson, K.A., 2016. Ultra-small inorganic nanoparticles: state-of-the-art and perspectives for biomedical applications. *Nanomed. Nanotechnol.* 12 (6), 1663–1701.
- Zeng, X., Ma, L.Q., Qiu, R., Tang, Y., 2009. Responses of non-protein thiols to Cd exposure in Cd hyperaccumulator *Arabis paniculata* Franch. *Environ. Exp. Bot.* 66, 242–248.
- Zeng, G., Wan, J., Huang, D., Hu, L., Huang, C., Cheng, M., Xue, W., Gong, X., Wang, R., Jiang, D., 2017. Precipitation, adsorption and rhizosphere effect: the mechanisms for phosphate-induced Pb immobilization in soils—a review. *J. Hazard. Mater.* 339, 354–367.
- Zhang, H., Ma, D., Qiu, R., Tang, Y., Du, C., 2017. Non-thermal plasma technology for organic contaminated soil remediation: a review. *Chem. Eng. J.* 313, 157–170.
- Zhang, Y.F., Sheng, J.C., Lu, Q.Y., 2004. Review on the soil remediation technologies. *Gansu Agric. Sci. Technol.* 10, 36–38.
- Zhang, F.Q., Zhang, H.X., Wang, G.P., Xu, L.L., Shen, Z.G., 2009. Cadmium induced accumulation of hydrogen peroxide in the leaf apoplast of *Phaseolus aureus* and *Vicia sativa* and the roles of different antioxidant enzymes. *J. Hazard. Mater.* 168, 76–84.
- Zhang, B.Y., Zheng, J.S., Sharp, R.G., 2010. Phytoremediation in engineered wetlands: mechanisms and applications. *Procedia Environ. Sci.* 2, 1315–1325.
- Zhao, L., Cao, X., Zheng, W., Scott, J.W., Sharma, B.K., Chen, X., 2016. Coprolysis of biomass with phosphate fertilizers to improve biochar carbon retention, slow nutrient release, and stabilize heavy metals in soil. *ACS Sustain. Chem. Eng.* 4 (3), 1630–1636.
- Zhao, C., Gupta, V.V., Degryse, F., McLaughlin, M.J., 2017. Abundance and diversity of sulphur-oxidising bacteria and their role in oxidising elemental sulphur in cropping soils. *Biol. Fertil. Soils* 53 (2), 159–169.
- Zhou, Q.X., Song, Y.F., 2004. Remediation of contaminated soils: principles and methods. *Sci. Press* 7, 345–346.
- Zhu, H., Chen, C., Xu, C., Zhu, Q., Huang, D., 2016a. Effects of soil acidification and liming on the phytoavailability of cadmium in paddy soils of central subtropical China. *Environ. Pollut.* 219, 99–106.
- Zhu, H., Chen, C., Xu, C., Zhu, Q., Huang, D., 2016b. Effects of soil acidification and liming on the phytoavailability of cadmium in paddy soils of central subtropical China. *Environ. Pollut.* 219, 99–106.
- Zhu, Y.L., Zayed, A.M., Quian, J.H., Souza, M.D.E., Terry, N., 1999. Phytoaccumulation of trace elements by wetland plants: II. Water hyacinth. *J. Environ. Qual.* 28, 339–344.
- Zhuang, P., Ye, Z.H., Lan, C.Y., Xie, Z.W., Shu, W.S., 2005. Chemically assisted phytoextraction of heavy metal contaminated soils using three plant species. *Plant Soil* 276 (1–2), 153–162.